

## WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6: WO 99/19492 (11) International Publication Number: A2 C12N 15/29, C07K 14/415, C12N 5/04, 22 April 1999 (22.04.99) (43) International Publication Date: 5/14, A01H 5/00, C12N 15/10, 15/82 (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, (21) International Application Number: PCT/EP98/06977 BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, 9 October 1998 (09.10.98) (22) International Filing Date:

ΑU

(71) Applicant (for all designated States except US): RHONE-POULENC AGRO [FR/FR]; 14/20, rue Pierre Baizet, F-69009 Lyon (FR).

10 October 1997 (10.10.97)

(72) Inventors; and

(30) Priority Data:

PO 9745

(75) Inventors/Applicants (for US only): DOUTRIAUX,
Marie-Pascale [FR/FR]; 64, route de Villebon, F-91160 Saulx les Chartreux (FR). BETZNER, Andreas, Stefan [AU/AU]; 40 Dallachy Place, Page, ACT 2614 (AU). FREYSSINET, Georges [FR/FR]; 21, rue de Nervieux, F-69450 Saint Cyr au Mont d'Or (FR). PEREZ, Pascal [FR/FR]; 17, chemin de la Pradelle, Varennes, F-63450 Chanonat (FR).

(74) Agent: GENIN, Patrick; Rhône-Poulenc Agro, DPI, 14/20, rue Pierre Baizet, F-69009 Lyon (FR).

KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published

Without international search report and to be republished upon receipt of that report.

(54) Title: METHODS FOR OBTAINING PLANT VARIETIES

(57) Abstract

An isolated and purified DNA molecule comprising a polynucleotide sequence encoding a polypeptide functionally involved in the DNA mismatch repair system of a plant.

#### FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Аппеліа	FI	Finland	LT	Lithuania	SK	Słovakia
AT	Austria	FR	Prance	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Мопасо	TD	Chad
BA	Bosnia and Herzegovina	CE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GII	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
ВG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	us	United States of America
CÁ	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
СН	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
C.W	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	L!	Liechtenstein	SD	Sudan		
DK	Denniark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	1.R	Liberia	SG	Singapore		

#### Methods for Obtaining Plant Varieties

#### TECHNICAL FIELD

The present invention relates to nucleotide sequences which encode polypeptides involved in the DNA mismatch repair systems of plants, and to the polypeptides encoded by those nucleotide sequences. The invention also relates to nucleotide sequences and polypeptide sequences for use in altering the DNA mismatch repair system in plants. The invention also relates to a process for altering the DNA mismatch repair system of a plant cell, to a process for increasing genetic variations in plants and to processes for obtaining plants having a desired characteristic.

#### BACKGROUND OF THE INVENTION

10

Plant breeding essentially relies on and makes use of genetic variation which occurs naturally within and between members of a family, a genus, a species or a subspecies. Another source of genetic variation is the introduction of genes from other organisms which may or may not be related to the host plant.

Allelic loci or non-allelic genes which constitute or contribute to desired quantitative (e.g. growth performance, yield, etc.) or qualitative (e.g. deposition, content and composition of seed storage products; pathogen resistance genes: etc.) traits that are absent, incomplete or inefficient in a species or subspecies of interest are typically introduced by the plant breeder from other species or subspecies, or *de novo*. This introduction is often done by crossing, provided that the species to be crossed are sexually compatible. Other means of introducing genomes, individual chromosomes or genes into plant cells or plants are well known in the art. They include cell fusion, chemically aided transfection (Schocher et al., 1986, Biotechnology 4: 1093) and ballistic (McCabe et al., 1988, Biotechnology 6: 923), microinjection (Neuhaus et al., 1987, TAG 75: 30), electroporation of protoplasts (Chupeau et al., 1989, Biotechnology 7: 53) or microbial transformation methods such as Agrobacterium mediated transformation (Horsch et al., 1985, Science 227: 1229; Hiei et al., 1996, Biotechnology 14: 745).

However, when a foreign genome, chromosome or gene is introduced into a plant, it will often segregate in subsequent generations from the genome of the recipient plant or plant cell during mitotic and meiotic cell divisions and, in consequence, become lost from the host plant or plant cell into which it had been introduced. Occasionally, however, the introduced genome, chromosome or gene physically combines entirely or in part with the genome, chromosome or gene of the host plant or plant cell in a process which is called recombination.

Recombination involves the exchange of covalent linkages between DNA molecules in regions of identical or similar sequence. It is referred to here as homologous recombination if donor and recipient DNA are identical or nearly identical (at least 99%).

hase sequence identity), and as homeologous recombination if donor and recipient DNA are not identical but are similar (less than 99% base sequence identity).

The ability of two genomes, chromosomes or genes to recombine is known to depend largely on the evolutionary relation between them and thus on the degree of sequence similarity between the two DNA molecules. Whereas homologous recombination is frequently observed during mitosis and meiosis, homeologous recombination is rarely or never seen.

From a breeder's perspective, the limits within which homologous recombination occurs, therefore, define a genetic barrier between species, varieties or lines, in contrast to homeologous recombination which can break this barrier. Homeologous recombination is thus of great importance for plant breeding. Accordingly there is a need for a process for enhancing the frequency of homeologous recombination in plants. In particular, there is a need for a process of increasing homeologous recombination to significantly shorten the length of breeding programs by reducing the number of crosses required to obtain an otherwise rare recombination event.

At least in Escherichia coli. homologous and homeologous recombination are known to share a common pathway that requires among others the proteins RecA, RecB, RecC. RecD and makes use of the SOS induced RuvA and RuvB, respectively. It has been suggested that mating induced recombination follows the Double-Strand Break Repair 20 model (Szostak et al., 1983, Cell 33, 25-35), which is widely used to describe genetic recombination in eukaryotes. Following the alignment of homologous or homeologous DNA double helices the RecA protein mediates an exchange of a single DNA strand from the donor helix to the aligned recipient DNA helix. The incoming strand screens the recipient helix for sequence complementarity, seeking to form a heteroduplex by hydrogen 25 bonding the complementary strand. The displaced homologous or homeologous strand of the recipient helix is guided into the donor helix where it base pairs with its counterpart strand to form a second heteroduplex. The resulting branch point then migrates along the aligned chromosomes thereby elongating and thus stabilising the initial heteroduplexes. Single stranded gaps (if present) are closed by DNA synthesis. The strand cross overs 30 (Holliday junction) are eventually resolved enzymatically to yield the recombination products.

Although in wild type E. coli homologous and homeologous recombination are thus mechanistically similar if not identical, homologous recombination in conjugational crosses E. coli x E. coli occurs five orders of magnitude more frequently than homeologous recombination in conjugational crosses E. coli x S. typhimurium (Matic et al. 1995; Cell 80, 507-515). The imbalance in favour of homologous recombination was shown to be caused largely by the bacterial MisMatch Repair (MMR) system since its

inactivation increased the frequency of homeologous recombination in E. coli up to 1000 fold (Rayssiguier et al. 1989, Nature 342, 396-401).

In E. coli, the MMR system (reviewed by Modrich 1991, Annual Rev Genetics 25, 229-253) is composed of only three proteins known as MutS, MutL and MutH. MutS recognizes and binds to base pair mismatches. MutL then forms a stable complex with mismatch bound MutS. This protein complex now activates the MutH intrinsic single stranded endonuclease which nicks the strand containing the misplaced base and thereby prepares the template for DNA repair enzymes.

During recombination, MMR components inhibit homeologous recombination. In vitro experiments demonstrated that MutS in complex with MutL binds to mismatches at the recombination branch point and physically blocks RecA mediated strand exchange and heteroduplex formation (Worth et al., 1994; PNAS 91, 3238-3241). Interestingly, the SOS dependent RuvAB mediated branch migration is insensitive to MutS/MutL, explaining the observed slight increase in SOS dependent homeologous recombination.

15 Homeologous mating even induces the SOS response, thereby taking advantage of RuvAB induction (Matic et al. 1995, Cell 80, 507-515).

The MMR system thus appears to be a genetic guardian over genome stability in E. coli. In this role it essentially determines the extent to which genetic isolation, that is, speciation, occurs. The diminished sensitivity of the SOS system to MMR, however, allows (within limits) for rapid genomic changes at times of stress, providing the means for fast adaptation to altered environmental conditions and thus contributing to intraspecies genetic variation and species evolution.

The important role of MMR in preserving genomic integrity has been established also in certain eukaryotes. In its efficiency, the human MMR, for example, may even counteract potential gene therapy tools such as triple-helix forming oligonucleotides including RNA-DNA hybrid molecules (Havre et al., 1993, J. Virology 67: 7234-7331; Wang et al., 1995, Mol. Cell. Biol. 15: 1759-1768; Kotani et al., 1996, Mol. Gen. Genetics 250: 626-634; Cole-Strauss et al., 1996, Science 273: 1387-1389). Such oligonucleotides are designed to introduce single base changes into selected DNA target sequences in order to inactivate for example cancer genes or to restore their normal function. The resulting base mismatches however are recognised by the mismatch repair system which then directs removal of the mismatched base, thereby reducing the efficiency of oligonucleotide induced site-specific mutagenesis.

To date, two families of related genes, homologous to the bacterial MutS and MutL genes have been identified or isolated in yeast and mammals (recent reviews by Arnheim and Shibata, 1997, Curr. Opinion Genet. Dev. 7, 364-370; Modrich and Lahue, 1996, Annual Rev. Biochem. 65, 101-153; Umar and Kunkel, 1996, Eur. J. Biochem. 238, 297-307). Biochemical and genetic analysis indicated that eukaryotic MutS homologs (MSH)

and MutL homologs (MLH. PMS), respectively, fulfil similar protein functions as their bacterial counterparts. Their relative abundance, however, could reflect different mismatch specificity and/or specialisation for different tissues or organelles or developmental processes such as mitotic versus meiotic recombination.

To date, six different genes homologous to *MutS* have been isolated in yeast (yMSH), and their homologs have been found in mouse (mMSH) and human (hMSH), respectively. Encoded proteins yMSH2, yMSH3 and yMSH6 appear to be the main MutS homologs involved in MMR during mitosis and meiosis in yeast, where the complementary proteins MSH3 and MSH6 alternatively associate with MSH2 to recognise different mismatch substrates (Masischky et al., 1996, Genes Dev. 10, 407-420). Similar protein interactions have been demonstrated for the human homologs hMSH2, hMSH3 and hMSH6 (Acharya et al., 1996, PNAS 93, 13629-13634).

MutL homologs (MLH and PMS), recently reviewed by Modrich and Lahue (1996, Annual Rev. Biochem. 65, 101-133) have so far been found in yeast (yMLH1 and yPMS1), mouse (mPMS2) and human (hMLH1, hPMS1 and hPMS2). The hPMS2 is a member of a family of at least 7 genes (Horii et al., 1994, Biochem. Biophys. Res. Commun. 204, 1257-1264) and its gene product is most closely related to yPMS1. Prolla et al. (1994, Science 265, 1091-1093) presented evidence for yPMS1 and yMLH1 to physically associate with each other and, together, to interact with the MutS homolog yMSH2 to form a ternary complex involved in mismatch substrate binding.

However, while medical interest in mismatch repair has prompted extensive research on MMR in bacteria, yeast and mammals, MMR genes have not been isolated from higher plants prior to the present invention and no attempts to adjust the plant MMR to plant breeding needs have been reported.

#### SUMMARY OF THE INVENTION

25

According to a first embodiment of the invention, there is provided an isolated and purified DNA molecule comprising a polynucleotide sequence encoding a polypeptide functionally involved in the DNA mismatch repair system of a plant. In one form of this embodiment, the invention provides an isolated and purified DNA molecule comprising a polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human. More particularly, the invention provides polynucleotide sequences encoding polypeptides which are homologous to the mismatch repair polypeptides MSH3 and MSH6 of Saccharomyces cerevisiae. Still more particularly, the invention provides the coding sequences of the genes AtMSH3 and AtMSH6 of Arabidopsis thaliana. as defined hereinbelow, and polynucleotide sequences encoding polypeptides which are homologous to polypeptides encoded by AtMSH3 and AtMSH6.

According to a second embodiment of the invention, there is provided an isolated and purified polypeptide functionally involved in the DNA mismatch repair system of a plant, for example a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human such as a polypeptide encoded by the genes AtMSH3 or AtMSH6 of Arabidopsis thaliana, as defined hereinbelow.

According to a third embodiment of the invention, there is provided an isolated and purified DNA molecule comprising a polynucleotide sequence selected from the group consisting of (i) a sequence encoding a polynucleotide which is capable of interfering with the expression of a plant polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human and thereby disabling said plant polynucleotide sequence; and (ii) a sequence encoding a polypeptide capable of disrupting the DNA mismatch repair system of a plant.

According to a fourth embodiment of the invention there is provided a chimeric gene comprising a DNA sequence selected from the group consisting of (i) a sequence encoding a polynucleotide which is capable of interfering with the expression of a plant polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human and thereby disabling said plant polynucleotide sequence, and (ii) a sequence encoding a polypeptide capable of disrupting the DNA mismatch repair system of a plant: together with at least one regulation element capable of functioning in a plant cell. Examples of such regulation elements include constitutive, inducible, tissue type specific and cell type specific promoters such as 35S, NOS, PR1a. AoPR1 and DMC1. Typically, a chimeric gene of the fourth embodiment will also include at least one terminator sequence, more typically exactly one terminator sequence.

In the third and fourth embodiments, said interference, by said polynucleotide sequence, with the expression of a plant polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair peptide of a yeast or a human typically occurs by hybridisation or by co-suppression.

According to a fifth embodiment of the invention there is provided a plasmid or vector comprising a chimeric gene of the fourth embodiment. A vector of the fifth 30 embodiment may be, for example, a viral vector or a bacterial vector.

According to a sixth embodiment of the invention, there is provided a plant cell stably transformed, transfected or electroporated with a plasmid or vector of the fifth embodiment.

According to seventh embodiment of the invention, there is provided a plant 35 comprising a cell of the sixth embodiment.

According to an eighth embodiment of the invention, there is provided a process for at least partially inactivating a DNA mismatch repair system of a plant cell, comprising

transforming or transfecting said plant cell with a DNA sequence of the third embodiment or a chimeric gene of the fourth embodiment or a plasmid or vector of the fifth embodiment, and causing said DNA sequence to express said polynucleotide or said polypeptide.

According to a ninth embodiment of the invention, there is provided a process for increasing genetic variation in a plant comprising obtaining a hybrid plant from a first plant and a second plant, or cells thereof, said first and second plants being genetically different; altering the mismatch repair system in said hybrid plant; permitting said hybrid plant to self-fertilise and produce offspring plants; and screening said offspring plants for plants in which homeologous recombination has occurred. For example, homeologous recombination may be evidenced by new genetic linkage of a desired characteristic trait or of a gene which contributes to a desired characteristic trait.

According to a tenth embodiment of the invention there is provided a process for obtaining a plant having a desired characteristic, comprising altering the mismatch repair system in a plant, cell or plurality of cells of a plant which does not have said desired characteristic, permitting mutations to persist in said cells to produce mutated plant cells, deriving plants from said mutated plant cells, and screening said plants for a plant having said desired characteristic.

In a preferred form of the ninth and tenth embodiments of the invention, the step of altering the mismatch repair system comprises introducing into said hybrid plant, plant, cell or cells a chimeric gene of the fourth embodiment and permitting the chimeric gene to express a polynucleotide which is capable of interfering with the expression of a plant polynucleotide sequence in a mismatch repair gene of the hybrid plant, plant, cell or cells, or a polypeptide capable of disrupting the DNA mismatch repair system of the hybrid plant or cells.

In other embodiments, the invention provides (a) an oligonucleotide capable of hybridising at 45°C under standard PCR conditions to a DNA molecule of the first embodiment; (b) an oligonucleotide capable of hybridising at 45°C under standard PCR conditions to the DNA of SEQ ID NO: 18 and (c) an oligonucleotide capable of hybridising at 45°C under standard PCR conditions to the DNA of SEQ ID NO:30; with the proviso that the oligonucleotide of (a), (b) and (c) is other than SEQ ID NO:1 or SEQ ID NO:2.

### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 provides a diagrammatic representation of the primer sequences used to 35 isolate AtMSH3.

Figure 2 is a plasmid map of clone 52, showing restriction enzyme cleavage sites in the 5' half of the full-length cDNA for AtMSH3.

Figure 3 is a plasmid map of clone 13, showing restriction enzyme cleavage sites in the 3' half of the full-length cDNA for AtMSH3.

Figure 4 is a sequence listing of the coding sequence of AtMSH3, together with a deduced sequence of the encoded polypeptide.

Figure 5 is a protein alignment of yeast (Saccharomyces cerevisiae) and Arabidopsis thaliana MSH3 protein.

Figure 6 provides a diagrammatic representation of the primer sequences used to isolate AIMSH6.

Figure 7 is a plasmid map of clone 43, showing restriction enzyme cleavage sites in the 5' half of the full-length cDNA for *AtMSH6*.

Figure 8 is a plasmid map of clone 62, showing restriction enzyme cleavage sites in the 3' half of the full-length cDNA for AtMSH6.

Figure 9 is a sequence listing of the coding sequence of AtMSH6, together with a deduced sequence of the encoded polypeptide.

Figure 10 is a protein alignment of yeast (Saccharomyces cerevisiae) and Arabidopsis thaliana MSH6 protein.

Figure 11 is a genomic sequence listing of AtMSH6.

Figure 12 is a plasmid map of plasmid pPF13.

Figure 13 is a plasmid map of plasmid pPF14.

Figure 14 is a plasmid map of plasmid pCW186.

Figure 15 is a plasmid map of plasmid pCW187.

Figure 16 is a plasmid map of plasmid pPF66.

Figure 17 is a plasmid map of plasmid pPF57.

Figure 18 is a diagrammatic representation of an antisense gene construction for use in homeologous meiotic recombination.

Figure 19 is a plasmid map of plasmid p3243.

## DETAILED DESCRIPTION OF THE INVENTION

The present invention is based on the inventors' discovery that there exist in higher plants genes which are homologous to MMR genes in E. coli, and to MMR genes in yeasts and humans.

Thus, the inventors have identified genes, herein designated AtMSH3 and AtMSH6, of the plant Arabidopsis thaliana which encode the proteins AtMSH3 and AtMSH6. These plant proteins are homologous to yMSH3 and yMSH6, respectively. The present inventors have isolated cDNAs encoding the proteins AtMSH3 and AtMSH6 and have isolated the complete gene encoding AtMSH6. Given the teaching herein, other genes (for example AtMSH2, and genes of other plants) may be obtained which are involved in DNA mismatch repair in plants, including other genes which are depoted polypeptides homologous to MMR proteins of yeasts or humans, such as genes which encode

polypeptides homologous to yeast MSH2, MLH1 or PMS2, or to human MLH1, PMS1 or PMS2. For example, given the teaching herein, genes of members of the *Brassicaceae* family or of other unrelated families, for example the *Poaceae*, the *Solanaceae*, the *Asteraceae*, the *Malvaceae*, the *Fabaceae*, the *Linaceae*, the *Canabinaceae*, the *Dauaceae* and the *Cucurbitaceae* family, and which encode polypeptides homologous to MMR proteins of yeasts or humans may be obtained.

Examples of plants whose genes encoding polypeptides homologous to MMR proteins of yeasts or humans may be obtained given the teaching herein include maize, wheat, oats, barley, rice, tomato, potato, tobacco, capsicum, sunflower, lettuce, artichoke, safflower, cotton, okra, beans of many kinds including soybean, peas, melon, squash, cucumber, oilseed rape, broccoli, cauliflower, cabbage, flax, hemp, hops and carrot.

Within the meaning of the present invention, a first polypeptide is defined as homologous to a second polypeptide if the amino acid sequence of the first polypeptide exhibits a similarity of at least 50% on the polypeptide level to the amino acid sequence of the second polypeptide.

A procedure which may be followed to obtain genes AtMSH3 and AtMSH6 is described in Example 1. Essentially the same technique may be applied to obtain other mismatch repair genes of Arabidopsis thaliana, and essentially the same technique as 20 exemplified herein may be applied to cDNA obtained by reverse transcription of RNA from other plants. Alternatively, given the sequence information disclosed herein, other degenerate oligonucleotide primers, especially oligonucleotides of the invention which are capable of hybridising at 45°C under standard PCR conditions (such as the conditions described in Example 1 using primers UPMU and DOMU) to AtMSH3 and/or AtMSH6 25 may be designed and obtained for use in isolating sequences of plant mismatch repair genes which are homologous to AtMSH3 or AtMSH6, from other plants. oligonucleotides of the invention which are capable of hybridising at 45°C under standard PCR conditions to plant mismatch repair genes of plants other than Arabidopsis thaliana also fall within the scope of the present invention and may be utilised to obtain mismatch Typically, such oligonucleotides are capable of 30 repair genes of still other plants. hybridising at 45°C under standard PCR conditions to a DNA molecule which encodes a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or a human. The temperature at which oligonucleotides of the invention hybridise to AtMSH3 and/or AtMSH6, or to plant mismatch repair genes of plants other than Arabidopsis thaliana, or 35 to DNA molecules which encode polypeptides which are homologous to a mismatch repair polypeptide of a yeast or a human may be higher than 45°C, for example at least 50°C, or at least 55°C, or at least 60°C or as high as 65°C.

The successful gene isolation disclosed herein demonstrates for the first time the existence of MMR in higher plants and indicates the presence of other plant MMR genes. For example, genes encoding the plant homologs of MSH1, MSH2, MSH4, MSH5, PMS1, PMS2 and MLH1 may be identified given the teaching herein. Such genes, as well as those specifically described herein, separately or in combination, are useful in manipulating the plant MMR for plant breeding purposes. Thus, for example, the plant MMR may be altered by including in a plant cell a polynucleotide sequence as defined herein above with reference to the third embodiment of the invention, and causing the polynucleotide sequence to express either a polynucleotide which disables a plant MMR gene, or a polypeptide which disrupts the plant's MMR system.

The DNA molecule of the third embodiment of the invention includes a polynucleotide sequence (herein referred to as a MMR altering gene) which may for example encode sense, antisense or ribozyme molecules characterised by sufficient base sequence similarity or complementarity to the gene to be altered to permit the antisense or 15 ribozyme molecule to hybridise with the plant MMR gene in vivo or to permit the sense molecule to participate in co-suppression. Alternatively, the MMR altering gene may encode a protein or proteins which interfere with the activity of a plant MMR protein and thus disrupt the plant's MMR system. For example, such encoded proteins may be antibodies or other proteins capable of interfering with MMR protein function, such as by 20 complexing with a protein functionally involved in plant MMR thereby disrupting the MMR of the plant. An example of such a protein is the MSH3 protein of Arabidopsis thaliana described herein or a protein of another plant which is homologous to the MSH3 protein of A. thaliana. For instance, overexpression of MSH3 in a plant cell causes MSH2 present in the cell to be substantially completely complexed, disrupting the 25 mismatch repair mechanism or mechanisms in the cell which are functionally dependent on the presence of a complex of MSH2 with MSH6. Similarly, mismatch repair mechanisms which depend on the presence of a complex of MSH2 and MSH3 may be disrupted by the overexpression of MSH6.

A chimeric gene of the fourth embodiment, incorporating a MMR altering gene, may be prepared by methods which are known in the art. Similarly, the MMR altering gene may be introduced into a plant cell, regenerating tissue or whole plant by techniques known in the art as being suitable for plant transformation, or by crossing. Known transformation techniques include Agrobacterium tumefaciens or A. rhizogenes mediated gene transfer, ballistic and chemical methods, and electroporation of protoplasts.

The MMR altering gene or genes are typically expressed from suitable promoters. Suitable promoters may direct constitutive expression, such as the 35S or the NOS promoter. Usually, however, the promoter will direct either inducible or tissue specific (e.g. callus; embryonic tissue; etc.), cell type specific (e.g. protoplasts; meiocytes. etc.) or developmental (e.g. embryo) expression of the altering gene or genes, in order for the

35

MMR system to function in tissue types or cell types, or at developmental stages of the plant, in which it is not desirable for the MMR system to be altered. Using such promoters, therefore, the activity of a MMR altering gene may be limited to a specific stage during plant development or it may be altered by controlling conditions external to the plant, and the deleterious effects of a permanently disabled or altered DNA mismatch repair system in a plant may be avoided. Examples of suitable promoters which are not constitutive are known in the art and include inducible promoters such as PR1a (reviewed by Gatz, 1997, Annual Rev. Plant Phys. Plant Mol. Biol. 48: 89), tissue specific promoters such as AoPR1 (Sabahattin et al., 1993, Biotechnology 11: 218), and cell-type specific promoters such as DMC1.

A chimeric gene in accordance with the invention may further be physically linked to one or more selection markers such as genes which confer phenotypic traits such as herbicide resistance, antibiotic resistance or disease resistance, or which confer some other recognisable trait such as male sterility, male fertility, grain size, colour, growth rate, flowering time, ripening time, etc.

The process of the tenth embodiment of the invention provides, for example, a process for generating intraspecies genetic variation by altering the mismatch repair system in a plant cell, in regenerating plant tissue or in a whole plant. The plant cell, regenerating tissue or whole plant includes and expresses one or more MMR altering 20 genes which are capable of altering mismatch repair in the plant cell, regenerating tissue or whole plant. Alteration of MMR may be achieved, for example, by inactivating the genes encoding plant MSH3 and/or plant MSH6. It is preferred to inactivate the plant MSH3 and MSH6 encoding genes at the same time and in the same plant cell, regenerating tissue or whole plant. Typically in this preferred form of the invention 25 inactivation of either plant MSH3 or MSH6 alone is insufficient to substantially alter the plant's mismatch repair system and only when both MSH3 and MSH6 are inactivated simultaneously is the plant's mismatch repair system sufficiently altered to prevent the MMR system from recognising base pair mismatches, base insertions or deletions as a result of DNA replication errors, DNA damage, or oligonucleotide induced site-specific 30 mutagenesis. However, in some applications of the invention, inactivation of only one gene may also be used to cause genomic instability or increase the efficiency of sitespecific mutagenesis.

If desired, the MMR altering gene or genes may later be rendered non-functional or ineffective, or may be removed from the genome of the plant cell, regenerating tissue or whole plant in order to restore mismatch repair in the plant cell, regenerating tissue or whole plant. The MMR altering gene or genes may be inactivated by means of known gene inactivation tools, such as ribozymes, or may be removed from the genome using gene elimination systems known in the art, such as CRE/LOX. It is preferred to render two genes, whose gene products combine to incapacitate MMR, ineffective by separating

the altering genes through segregation. Therefore, in a preferred embodiment of the invention a first plant cell or plant is generated in which only plant MSH3 is incapacitated, and a second plant cell or plant is generated in which only plant MSH6 is incapacitated. The combination of both genomes, for example by crossing, then produces significant MMR deficiency in those cells or plants which have inherited both altering genes. If the altering genes are expressed from unlinked loci, gene segregation restores MMR activity in the progeny of the cells or plants.

In a process of the ninth embodiment of this invention, homeologous recombination is enhanced between different genomes, chromosomes or genes in plant cells or plants by altering MMR in said plant cells or plants. Such genomes, chromosomes or genes are characterised in that they originate from different plant families, genera, species, subspecies, plant varieties or lines. Hybrid plant cells or hybrid plants may be produced by crossing, by cell fusion or by other techniques known in the art. These plant cells or plants are further characterised by expressing one or more genes that are capable of altering mismatch repair in the plant cell or plants.

In the process of the ninth embodiment, the homeologous recombination is typically for the purpose of introducing a desired characteristic in the hybrid plant. In this typical application of the process of the ninth embodiment, and in the process of the tenth embodiment the desired characteristic may be any characteristic which is of value to the plant breeder. Examples of such characteristics are well known in the art and include altered composition or quality of leaf or seed derived storage products (e.g. oil, starch, protein), altered composition or quality of cell walls (e.g. decrease in lignin content), altered growth rate, prolonged flowering, increased plant yield or grain yield, altered plant morphology, resistence to pathogens, tolerance to or improved performance under environmental stresses of various kinds, etc.

In a preferred form of the tenth embodiment, an MMR altering gene is cointroduced along with the homeologous genome, chromosome or gene of another plant
cell or plant into an MMR proficient plant cell or MMR proficient plant to produce a
hybrid plant cell or hybrid plant in which homeologous recombination can occur.

Suitably, the MMR proficient plant cell or MMR proficient plant may also include an
MMR altering gene. For example a gene capable of inactivating plant MSH3 may be cointroduced along with the homeologous genome, chromosome or gene of another plant
cell or plant into an MMR proficient plant cell or MMR proficient plant in which MSH6
is inactivated. A resultant hybrid plant in which homeologous recombination occurs will
include both the MSH3 and MSH6 altering genes and its MMR system will therefore be
inactivated.

In this form of the invention, if hybrid plants are to be produced by crossing, the MMR altering gene preferably originates from the male parent, thus ensuring that the

MMR altering gene is always introduced and is not present in the recipient cell. That is, the MMR of the recipient cell, prior to introduction of the MMR altering gene, is typically proficient. Alternatively, if an MMR altering gene is present in a recipient cell it may be ineffective or inefficient on its own, or it may be linked to an inducible or tissue specific or cell type specific promoter which only renders the MMR altering gene active under limited conditions.

Thus, in a preferred form of the process of the ninth embodiment, the MMR system of the hybrid plant is initially unaltered. In this form of the process, the step of altering the mismatch repair system may comprise introducing into the hybrid plant, or cells thereof, a MMR altering gene, such as by Agrobacterium tumefaciens or A. rhizogenes mediated gene transfer, ballistic and chemical methods, and electroporation of protoplasts.

The MMR altering gene or genes are typically expressed from suitable promoters, as described above. Preferably, the promoter is transcriptionally active in mitotically and meiotically active tissue and/or cells to ensure MMR alteration after chromosome pairing at mitosis and meiosis, respectively. The preferred timing for MMR alteration is at meiosis, because recombinant genomes, chromosomes or genes are directly transmitted to the progeny. A suitable meiocyte specific promoter is for example the *DMC*1 promoter from *Arabidopsis thaliana* ssp. *Ler.* (Klimyuk and Jones, 1997, Plant J. 11, 1-14). However, mitotic homeologous recombination is also a desirable outcome as somatic recombination events can be transmitted to offspring due to the totipotency of plant cells and the lack of predetermined germ cells in plants.

If desired, the MMR altering gene or genes may later be rendered non-functional or ineffective, or may be removed from the hybrid plant or hybrid plant cells, in order to restore mismatch repair in the hybrid plant or hybrid plant cells. The MMR altering gene or genes may be inactivated by means of known gene inactivation tools as described herein above.

#### **EXAMPLES**

## Example 1. Cloning of the AtMSH3 and AtMSH6 coding sequences

Isolation of partial AtMSH3 and AtMSH6 consensus sequences

Degenerate oligonucleotides UPMU (SEQ ID NO:1) and DOMU (SEQ ID NO:2)

UPMU CTGGATCCACIGGICCIAA(C/T)ATG

DOMU CTGGATCC(A/G)TA(A/G)TGIGTI(A/G)C(A/G)AA

were used to isolate AtMSH3 and AtMSH6 sequences by PCR amplification.

Primers UPMU and DOMU correspond to conserved amino acid sequences of the proteins MutS (E. coli and S. typhimurium), HexA (S. pneumoniae). Rep1 (mouse) and Duc1 (human). The conserved regions to which they are targeted are TGPNM for UPMU (amino acid positions 852-856 for AtMSH6 and 816-820 for AtMSH3) FATHY or FVTHY

for DOMU (amino acid positions 964-968 for AtMSH6 and 928-932 for AtMSH3, respectively.) These primers have been used to isolate MSH2 and MSH1 from yeast (Reenan and Kolodner, Genetics 132: 963-973 (1992)) and MSH2 from *Xenopus* and mouse (Varlet et al., Nuc. Acids Res. 22:5723-5728 (1994)).

Template single strand cDNA was produced by reverse transcription of 2 µg total RNA from a cell suspension culture of Arabidopsis thaliana ecotype Columbia (Axelos et al. 1989, Mol. Gen. Genetics 219: 106-112). The PCR reaction was performed under the following conditions in a final volume of 100µl: 0.2mM dNTP, 1µM each primer, 1XPCR buffer, lu Taq DNA polymerase (Appligene) in the presence of template cDNA. PCR 10 parameters were 5 minutes at 94°C, followed by 30 cycles of 40 seconds at 95°C, 90 seconds at 45°C, 1 minute at 72°C. The amplification products were cloned into pGEM-T vector (Promega) and sequenced. Two different clones were isolated, S5 (350bp) was homologous to MSH3, S8 (327bp) was homologous to MSH6. Complete cDNA sequences were then isolated according to the Marathon cDNA amplification kit procedure (Clontech). 15 In summary, this procedure involves producing double stranded cDNA by reverse transcription of 2µg polyA+ RNA from the cell suspension culture of Arabidopsis. Adaptors are ligated on each side of the cDNA. The ligated cDNA is used as a template for 5' and 3' RACE PCR reactions in the presence of primers that are specific for the adaptor on one side (AP1 and AP2), and specific for the targeted gene on the other side. A 5' and a 3' 20 fragment that overlap are thus produced for each gene. The complete gene coding sequence can be reconstituted taking advantage of a unique restriction site, if available, in the overlapping region. Specific details of this procedure as it was used to isolate AtMSH3 and AIMSH6 coding regions, are as follows.

### Isolation of AIMSH3 complete coding sequence

25

From the sequence of clone S5, primer 636 (SEQ ID NO:3) was designed:

636 TGCTAGTGCCTCTTGCAAGCTCAT.

Primer AP1 (SEQ ID NO:4) is complementary to a portion of an adaptor sequence which had been ligated to the 5' and 3' ends of Arabidopsis cDNA:

API CCATCCTAATACGACTCACTATAGGGC.

PCR performed on the ligated cDNA with primers 636 and AP1 for the 5' RACE PCR was followed by a second round of amplification with the nested primers AP2 (SEQ ID NO:5) and S525 (SEQ ID NO:6)

AP2 ACTCACTATAGGGCTCGAGCGGC

S525 AGGTTCTGATTATGTGTGACGCTTTACTTA

35 (the latter was also designed to correspond to a part of the sequence of clone S5) and produced a 2720bp DNA fragment. Figure 1 provides a diagrammatic representation of the primer sequences used to isolate AIMSH3. Another primer (S51, SEQ ID NO:7)

S51 GGATCGGGTACTGGGTTTTGAGTGTGAGG

was designed closer to the 5° border and permitted the determination of 99bp upstream to the ATG initiation codon. For the 3° RACE PCR, a first PCR reaction was performed with primers AP1 and 635 (SEQ ID NO:8).

- 635 GCACGTGCTTGATGGTGTTTTCAC
- 5 followed by a second round of amplification, using the nested primers AP2 and S523 (SEQ ID NO:9)
  - S523 TCAGACAGTATCCAGCATGGCAGAAGTA

which produced a DNA fragment of 890bp. Both DNA fragments were subcloned into pGEM-T and sequenced. Since PCR amplification using the Expand Long Template PCR system (Boehringer-Mannheim) produced errors in the sequence, new oligonucleotides were designed to isolate those sequences again by PCR, but with the high fidelity DNA polymerase Pfu. PCR with primers 1S5 (SEQ ID NO:10) and S53 (SEQ ID NO:11)

- 1S5 ATCCCGGGATGGGCAAGCAAAAGCAGCAGACGA
- S53 GACAAAGAGCGAAATGAGGCCCCTTGG
- amplified the 1244bp fragment clone 52 (SEQ ID NO:12, cloned into pUC18/Sma1). PCR with primers S52 (SEQ ID NO:13) and 2S5 (SEQ ID NO:14)
  - 2S5 ATCCCGGGTCAAAATGAACAAGTTGGTTTTAGTC
  - S52 GCCACATCTGACTGTTCAAGCCCTCGC

amplified the 2104bp clone 13 (SEQ ID NO:15, cloned into pUC18/Sma1). The complete coding sequence of the AtMSH3 gene was reconstructed in pUC18 by ligating the 5' half of AtMSH3 (clone 52) to the 3' half of AtMSH3 (clone 13) after digesting with BamH1 which has a unique cleavage site in the overlapping region of both clones. This manipulation yielded plasmid pPF26. The Smal fragment from pPF26 contains the complete AtMSH3 coding sequence. The remaining primers referred to in Figure 1 are as follows:

- S51 GGATCGGGTACTGGGTTTTGAGTGTGAGG (SEQ ID NO:16)
- S525 AGGTTCTGATTATGTGTGACGCTTTACTTA (SEQ ID NO:17)

Figures 2 and 3 provide plasmid maps of clones 52 and 13 respectively, showing restriction enzyme cleavage sites. The complete AtMSH3 coding sequence (SEQ ID NO:18) 30 is 3246bp long and is shown in Figure 4 together with the deduced sequence (SEQ ID NO:19) of the encoded polypeptide. AtMSH3 is clearly homologous to the yeast and mouse MSH3 genes. A sequence alignment of polypeptides encoded by AtMSH3 and that encoded by Saccharomyces cerevisiae MSH3 is set out in Figure 5.

#### Isolation of the AtMSH6 complete coding sequence and genomic sequences

- The same procedure allowed isolation of the AtMSH6 cDNA. Figure 6 provides a diagrammatic representation of the primer sequences used to isolate AtMSH6. For the 5' RACE PCR, primers 638 (SEQ ID NO:20) and AP1 (SEQ ID NO:4)
  - 638 TCTCTACCAGGTGACGAAAAACCG allowed the amplification of a 2889 DNA fragment. Primer S81 (SEQ ID NO:21)

## S81 CGTCGCCTTTAGCATCCCCTTCCTTCAC

helped define the 142bp upstream to the ATG initiation codon. On the 3' side, RACE PCR was initially performed with primers S823 (SEQ ID NO:22) and API (SEQ ID NO:4),

S823 GCTTGGCGCATCTAATAGAATCATGACAGG

5 and then with the nested primers 637 (SEQ ID NO:23) and AP2 (SEQ ID NO:5).

637 GACAGCGTCAGTTCTTCAGAATGC

to produce a 774bp DNA fragment. As for AtMSH3, those fragments were cloned and sequenced. Re-isolation of the DNA sequence using the high fidelity Pfu polymerase and newly designed primers 1S8 (SEQ ID NO:24) and S83 (SEQ ID NO:25) (for the 5' side) led to a 2182 bp DNA fragment identified as clone 43 (SEQ ID NO:26, cloned in pUC18/Sma1), and a 1379bp clone identified as clone 62 (SEQ ID NO:27, also cloned in pUC18/Sma1).

1S8 ATCCCGGGATGCAGCGCCAGAGATCGATTTTGT

2S8 ATCCCGGGTTATTTGGGAACACAGTAAGAGGATT (SEQ ID

NO:28)

15

S82 GCGTTCGATCATCAGCCTCTGTGTTGC (SEQ ID NO:29)

S83 CGCTATCTATGGCTGCTTCGAATTGAG

Figures 7 and 8 provide plasmid maps of clones 43 and 62 respectively, showing restriction enzyme cleavage sites. Clones 43 and 62 were digested by the *Xmn1* restriction enzyme for which a unique site is present in their overlapping region and then ligated. The complete *AtMSH6* coding sequence (SEQ ID NO:30) is 3330bp long and is shown in Figure 9 together with the deduced sequence (SEQ ID NO:31) of the encoded polypeptide. *AtMSH6* is clearly homologous to the yeast and mouse *MSH6*genes. A sequence alignment of polypeptides encoded by *AtMSH6* and that encoded by *Saccharomyces cerevisiae MSH6* is set out in Figure 10.

An AtMSH6 genomic sequence was also isolated from a genomic DNA library constituted after partial Sau3AI digestion of DNA from the Arabidopsis cell suspension. 8062bp were sequenced that covered the AtMSH6 gene and show colinearity with the cDNA. 16 introns are found scattered along the gene. The complete genomic sequence 30 (SEQ ID NO:98) is shown in Figure 11.

## Example 2. A measure of somatic variation in MMR deficient plants

#### Constructs

Constructs with antisense AtMSH3 or antisense AtMSH6 or both AtMSH3/AtMSH6 under the control of a single 35S promoter have been inserted into the binary vector pPZP121 (Hajdukiewicz et al., Plant Mol. Biol. 23, 793-799) between the right and left borders of the T-DNA. The pPZP121 plasmid confers chloramphenical resistance to Escherichia coli or Agrobacterium tumefaciens bacteria. The uacC1 gene is carried by the T-DNA and allows selection of transformed plant cells on gentamycin Majdukiewicz et al., Plant Mol. Biol. 25, 989-994). For the purpose of expressing antisense constructs, a 35S

promoter/terminator cassette with a polylinker was introduced into pPZP121. The 3' ends of the considered genes have been chosen since this region seems more efficient for antisense inhibition. For AtMSH3 this corresponds to clone 13 (2104bp), for AtMSH6 this corresponds to clone 62 (1379bp). Clone 13 comprises 2104bp of the 3' region that were cut off the pUC18 vector by Sal1/Sst1 restriction, blunted with T4 DNA polymerase and ligated into the T4 DNA polymerase blunted BamHI site of pPZP121/35S, creating clone pPF13. The same procedure was followed for the 3' region of AtMSH6 clone 62 (1379bp) thus creating plasmid pPF14. For the double constructs, the 3' region (from clone 62) of AtMSH6 was introduced ahead of the AtMSH3 region into pPF13 creating pCW186 and reciprocally, the 3' region of AtMSH3 (from clone 13) was introduced ahead of AtMSH6 into pPF14, creating pCW187.

These constructs were introduced into the Arabidopsis cells (as described below) of wildtype Columbia and of the Columbia tester line.

An alternative strategy to antisense inhibition of AtMSH6 comes from experiments of Marra et al. (1998. Proc. Natl. Acad. Sci USA 95. 8568-8573) who show that overexpression of functional MSH3 results in depletion of MSH6 protein in human cells. This depletion may generate a mismatch repair mutant phenotype.

For the purpose of overexpressing functional AtMSH3 protein in plant cells, the complete MSH3 coding region was excised from pPF26 (example 1) by digestion with 20 SmaI, and was inserted into the SmaI site of pCW164. The resulting construct was named pPF66. It contains a complete AtMSH3 gene under the control of the 35S promoter inside the left (LB) and right (RB) border of the T-DNA. This T-DNA also contains the hpt2 gene for gentamycin selection. Plasmid pPF66 was introduced into Arabidopsis cells as described below. One cell clone was selected which clearly overexpressed the AtMSH3 gene as shown by Northern analysis. Figures 12-16 provide plasmid maps of plasmids pPF13, pPF14, pCW186, pCW187 and pPF66, respectively.

#### Construction of tester construct

For the purpose of Forward Mutagenesis Assays, a tester construct was built containing the coding regions for nptII, codA, uidA. All three genes are driven by the 35S promoter and are terminated by the 35S terminator. This construct was obtained by introducing an EcoR1 fragment encoding the codA cassette (2.5kb) and a HindIII fragment encoding the uidA (GUS) cassette (2.4kb) into the pPZP111 vector (Hajdukiewicz et al.,1994, Plant Mol Biol 23: 793-799) which already contained the nptII expression cassette. This new plasmid was named pPF57. NptII is used to select for transformed plant cells.

35 GUS is used to analyse the degree of gene silencing in the construct (i.e. to identify cell lines in which the transgenes are expressed), and codA is used as a marker for forward mutagenesis (described below).

The plasmid map of pPF57 is provided in Figure 17.

#### Plant cell transformation

The constructs are introduced into Agrobucterium by electroporation. Plant cells are then transformed by co-cultivation. A suspension culture of Arabidopsis thaliana cells that 5 has been established by Axelos et al. (1992, Plant Physiol. Biochem. 30, 1-6) may be used. One day old freshly subcultured cells are diluted five times into AT medium (Gamborg B5 medium. 30g/l sucrose, 200µg/l NAA). 10µl of saturated Agrobacterium containing the transforming T-DNA constructs are added to 10ml diluted cells in a 100ml erlenmeyer. The co-cultivation is agitated slowly (80rpm) for 2 days. The cells are then washed 3 times into AT medium and finally resuspended in the same initial volume (10ml). The culture is agitated for 3 days to allow expression before plating on selection plates (AT/BactoAgar 8g/l+gentamycin 50µg/ml). Transformed individual calli are isolated 3 weeks later.

#### Tester Strain

The tester construct on plasmid pPF57 was introduced into Arabidopsis cells of wildtype Columbia using the transformation protocol described above. Among 10 candidate transformants, one cell clone was shown (by Southern analysis) to have a unique T-DNA insertion. All three genes were shown to be functional in this cell line as indicated by resistance to kanamycin, blue staining in the presence of X-Glu (GUS), and sensitivity to 5-fluoro-cytosine (codA).

MMR altering genes (described above) were then introduced individually into the tester line and transformed cells are used for analysis of both Microsatellite Instability and Forward Mutagenesis.

#### Microsatellite analysis

Microsatellites have been described in Arabidopsis (Bell and Ecker, 1994, Genomics 19, 137-144). The present Example is based on a study of instability of microsatellites that are polymorphic amongst different ecotypes. DNA is extracted from the transformed calli. Specific primers have been defined that are used to amplify the microsatellite sequence. One of the two primers is previously P<sup>32</sup> labelled by T4 kinase. In case of a polymorphic variation, new PCR products appear that do not follow the expected pattern of migration on a polyacrylamide gel. This is a commonly observed feature for MMR deficient cells in yeast or mammalian cells.

In particular, the present Example describes a study on microsatellites ca72 (CA<sub>18</sub>), nga172 (GA<sub>29</sub>), and ATHGENEA(A<sub>39</sub>), chosen because they belong to the types predominantly affected in human mismatch repair deficient tumors. The size of these microsatellites is not conserved from one *Arabidopsis* ecotype to the other.

Arabidopsis cells which are transformed with an MMR altering gene (above) and control cells not expressing the MMR altering gene are allowed to form calli. DNA is

rapidly extracted from the calli and is analysed for microsatellite instability as described in detail by Bell and Ecker 1994. Genomics 19, 137-144. In summary, the relevant microsatellite is amplified by PCR using P32 labelled primers. The PCR products are separated on a DNA sequencing gel for size determination. Size differences between microsatellites from transformed and control cells not expressing the MMR altering gene in question indicate microsatellite instability as a result of MMR alteration.

The sequences of primers used for PCR amplification of microsatellites ca72 and nga172 are included in Table 1. PCR amplification of microsatellite ATHGENEA made use of a forward primer containing the sequence

#### ACCATGCATAGCTTAAACTTCTTG (SEQ ID NO:32)

and of a reverse primer containing the sequence

#### ACATAACCACAAATAGGGGTGC (SEQ ID NO:33).

The amplification for microsatellite ca72 revealed in Columbia control cells (with respect to the MMR altering gene) a 248 bp long PCR fragment instead of the published length of 124 bp. DNA sequencing verified this fragment as a CA<sub>18</sub> microsatellite.

#### Forward mutagenesis assay

10

Tester cells transformed with antisense AtMSH3 or antisense AtMSH6 or both AtMSH3/AtMSH6 are analysed for the stability of the codA gene. The functional codA gene confers to sensitivity to 5-fluoro-cytosine (5FC), whereas a gene inactivating mutation in codA will confer resistance to 5FC. The frequency of resistant cells is therefore a good indicator of somatic variation as a direct result of MMR alteration. Variants resistant to 5FC are first analysed for GUS activity. If GUS is inactive, 5FC resistance is assumed to be due to gene silencing (all three genes are under the 35S promoter). If GUS is active, 5FC resistance is assumed to be due to forward mutations that have inactivated codA. PCR is then performed on the putative codA mutant genes which is then sequenced to confirm the presence of forward mutations in codA.

Besides codA, other marker genes may also be used for the Forward Mutagenesis Assay such as the ALS gene (conferring sensitivity to valine or to sulfonylurea; Hervieu and Vaucheret, 1996, Mol. Gen. Genet. 251 220-224; Mazur et al. 1987, Plant Physiol. 85 1110-30 1117).

## Example 3. Homeologous meiotic recombination in Arabidopsis thaliana

- A. Construction of a meiocyte specific gene expression cassette comprising the *DMC1* promoter and the *NOS* terminator
- (i) The DMC1 promoter may be used as published by Klimyuk and Jones, 1997, 35 Plant J. 11.1-14). To obtain a more convenient alternative for gene cloning, a 3.3 Kb

20

long subfragment of the DMC1 promoter was obtained by PCR from genomic DNA of Arabidopsis thaliana (ssp. Landsberg erecta "Ler").

The PCR was done in three rounds:

Round One: A 3.7 Kb long product was obtained using the forward primer 5 DMCIN-A comprising the sequence

GAAGCGATATTGTTCGTG (SEQ ID NO:34)

and the reverse primer DMCIN-B comprising the sequence

AGATTGCGAGAACATTCC (SEQ ID NO:35).

The weak amplification product was then used as template for round two and three.

Round Two: A 3.1 Kb long product comprising the promoter and the 5' untranslated leader was obtained using forward primer DMCIN-1, which contained the sequence

acgcgtcgacTCAGCTATGAGATTACTCGTG (SEQ ID NO:36)

and introduced a SaII cloning site at the 5' end of the promoter fragment, and reverse primer DMCIN-2 which contained the sequence

gctctagaTTTCTCGCTCTAAGACTCTCT (SEQ ID NO:37)

and introduced a XbaI site at the 3' end of the PCR fragment.

Round Three: A 0.2 Kb long product comprising the first exon/intron of the *DMC*1 promoter was obtained using forward primer DMCIN-3, which contained the sequence

gctctagaGCTTCTCTTAAGTAAGTGATTGAT (SEQ ID NO:38)

and introduced a XbaI site at the 5' end of the PCR fragment, and reverse primer DMCIN-4, containing the sequence

tecceeggetegagagatetecatggTTTCTTCAGCTCTATGAATCC (SEQ ID NO:39) and introduced at the 3' end of the PCR product restriction sites for Ncol, BgIII, Xhol and 25 Smal.

The products obtained in round Two and Three were digested with XbaI and subsequently ligated to reconstitute a 3.3 Kb long DMC1 promoter from which the first two in-frame ATG start codons were replaced with a unique restriction site for XbaI. This promoter can be cloned between the restriction sites for SaII and SmaI of p3264, which contains the SacI-EcoRI NOS terminator in pBIN19, to yield the entire expression cassette in pBIN19. This cassette contains the following cloning sites: NcoI, BgIII, XhoI. SmaI and (already present on p3264) KpnI and SacI.

(ii) Another strategy yielded the following convenient *DMC*1 promoter. A 1.8 kb DNA fragment comprising the 3' terminal part of the meiocyte specific *DMC*1 promoter was isolated by PCR from purified genomic DNA of Arabidopsis thaliana (ssp. Landsberg erecta "Ler"). The forward PCR primer (DMC1a) contained the sequence

acgcgtcgacGAATTCGCAAGTGGGG (SEQ ID NO:40)

and introduced a SaII cloning site at the 5' end of the promoter fragment. The reverse PCR primer (DMC1b) contained the sequence

tecatggagatetecegggtacCGATTTGCTTCGAGGG (SEQ ID NO:41) introducing a polylinker region at the 3' end of the promoter fragment. The PCR reaction was carried out with VENT DNA Polymerase (NEB) over 25 cycles using the following cycling protocol: 1 minute at 94°C, 1 minute at 54°C, 2 minutes at 72°C.

The PCR reaction yielded a blunt ended DNA fragment which was digested with restriction enzyme Sall and was cloned into the cleavage sites of restriction enzymes Sall and Smal in plasmid p2030, a pUC118 derivative containing the SacI-EcoRI NOS terminator fragment of pBIN121. The cloning yielded plasmid p2031, containing the DMC1 promoter-polylinker-NOS terminator expression cassette depicted in Figure 18.

#### 10 B. Construction of an MSH3 antisense gene under the control of the DMC1 promoter

A 2.1 kb DNA fragment encoding the carboxyterminal part of AtMSH3 was removed from the polylinker of clone 13 described in Example 1 after (i) digestion with KpnI. (ii) blunting of the DNA ends generated by KpnI and (iii) digestion with BamHI. The isolated fragment was then cloned in antisense orientation downstream of the DMC1 promoter in plasmid p2031, which had been digested with restriction enzymes SmaI and BglII. This cloning yielded plasmid p2033 (Figure 18).

After digestion of p2033 with EcoRI, a 4.1 kb DNA fragment was recovered comprising the DMC1 promoter, the partial MSH3 cDNA in antisense orientation with respect to the promoter and the NOS terminator. This fragment was cloned into the EcoRI restriction site of plant transformation vector pNOS-Hyg-SCV to yield plasmid p3242 (Figure 18).

## C. Construction of a combined MSH6/MSH3 antisense gene under the control of a single DMC1 promoter

A 3.1 kb fragment, encoding in antisense orientation the partial AtMSH6 and AtMSH3 sequences and the 35S terminator, was isolated from pCW186 by digestion with KpnI. The fragment was treated with Klenow enzyme to blunt both ends. It was then cloned into the SmaI site of plasmid p3243 (a pNOS-Hyg-SCV derivative, illustrated in Figure 19), which had been opened to delete the region between the SmaI sites. Clones containing the fragment in the antisense orientation with respect to the DMC1 promoter (described in A(ii) above) were identified by diagnostic digestion with BamHI. The selected construct was named p3261.

Another practical way of cloning the double antisense gene is as follows. A 1 kb DNA fragment encoding the carboxyterminal part of AtMSH6 is isolated from clone 62 described in Example 1 after digestion of clone 62 plasmid DNA with BamHI, which cleaves in the 5' polylinker region flanking the partial cDNA, and with EcoRI, which cleaves within the cDNA. The isolated fragment is treated with Klenow enzyme to blunt both its ends and is cloned into the recipient plasmid p2033 or p3242. For the purpose of

cloning, the recipient plasmid may be cleaved with either Aval or Ncol and can be blunted with Klenow enzyme to produce blunt acceptor ends for fragment cloning. This cloning yields two opposite orientations of cloned fragment DNA with respect to the DMC1 promoter. These can be identified by diagnostic digestion with restriction enzymes Scal or XmnI in conjunction with Sacl. The selected construct contains the DMC1 promoter, the combined partial cDNAs for AtMSH3 and AtMSH6 (both cloned in antisense orientation with respect to the DMC1 promoter) and the NOS terminator. If the recipient plasmid is p2033, the combined antisense gene under control the single DMC1 promoter is recovered from the vector after EcoRI digestion and is cloned into the EcoRI restriction site of pNOS-Hyg-SCV.

## D. Construction of a full-length MSH3 sense gene under control of the DMC1 promoter for overexpression of functional MSH3 protein

Overexpression of MSH3 protein was shown in human cells (Marra et al., 1998, Proc. Natl. Acad. Sci. USA 95, 8568-8573) to complex all available MSH2 protein. This leaves MSH6 protein without its partner, leading to the degradation of MSH6 protein and eventually to a mismatch repair phenotype.

This phenomenon is exploited to increase homeologous meiotic recombination in Arabidopsis as an alternative to antisense inhibition of AtMSH6. For this purpose the full-length cDNA encoding AtMSH3 is isolated from plasmid pPF66 by digestion with Smal and is cloned into the Smal site of the DMC1 expression cassettes described in A(i).

# E. Selection of Recombination markers on homeologous chromosomes of Arabidopsis thaliana subspecies Landsberg erecta (Ler), Columbia (Col) and C24, respectively

## E(i). Visual recombination markers in Arabidopsis th. subspecies C24:

Agrobacterium mediated transformation with a T-DNA containing a 35S-GUS gene, inactivated by insertion of a 35S-Ac transposable element (Finnegan et al., 1993, Plant Mol. Biol. 22, 625-633), had yielded a C24 line in which the T-DNA construct was integrated into chromosome 2. Genetic and molecular analysis of this line shows that the Ac transposon had excised from its T-DNA locus thereby restoring GUS activity, but had re-inserted into the chromosome at a distance of 16.4 cM, where it stayed fixed (due to disablement of Ac) within the chlorina gene. Insertional inactivation of the chlorina gene caused a bleached phenotype in those plants that were homozygous for this mutation. Because of the two linked phenotypic markers, chlorina3A:Ac and GUS, this C24 line was used in crosses to wildtype Ler for analysis of meiotic homeologous recombination on chromosome 2 in conjunction with molecular recombination markers.

### 35 <u>E(ii)</u>. Visual recombination markers in Arabidopsis th. Ler:

The Ler line NW1 (obtained from NASC, Nottingham, UK) contains one recessive visual marker per chromosome, i.e. an-1 on Chr.1, py-1 on Chr.2, gl1-1 on Chr.3, cer2-1

on Chr.4, and ms1-1 on Chr.5. This line is used in crosses to wildtype C24 which expresses an MMR altering gene for analysis of meiotic homeologous recombination on chromosomes 1-5 in conjunction with molecular recombination markers listed in Table 1.

Other Ler lines from NASC have several visual markers in close proximity to each other on the same chromosome. When these lines are used for hybrid production, analysis of homeologous meiotic recombination may make use entirely of visual recombination markers. Particularly suitable for crossing to C24 wildtype that is expressing a MMR altering gene are the following Ler lines:

NW22: relative markers are dis1 - (4 cM) - ga4 - (11 cM) - th1 on chromosome 1.

NW10: relevant markers are tz-201 - (5 cM) - cer3 on chromosome 5.

10

NW134, relevant markers are ttg - (4 cM) - ga3 on chromosome 5.

NW24 (abi3-1) and NW64 (gl1-1). When present in the same plant on chromosome 3, abi3-1 and gl1-1 are 13 cM apart. Since this marker combination is not available from NASC, we have combined these markers by crossing line NW24 to line NW64. The F1 offspring were allowed to self-fertilise and to produce F2 seeds. F2 Plants which carry both markers as homozygous traits on the same chromosome can be identified firstly, by germinating F2 seeds on germination medium containing selective concentrations of abscisic acid, and subsequently, by identifying among the abscisic acid resistant plants those individuals which show the glabra phenotype.

#### 20 E(iii) Molecular recombination markers in Col, Ler and C24:

The genome of Arabidopsis thaliana is interspersed with unique base sequences arranged as simple tandem repeats. Allelic repeats can vary in length between different Arabidopsis subspecies and when amplified by PCR yield diagnostic DNA products of different length named Simple Sequence Length Polymorphisms (SSLPs). Many SSLPs have been genetically mapped and have been assigned to unique chromosome locations on the recombinant inbred map (Bell and Ecker, 1994, Genomics 19, 137-144; Lister and Deans lines, Weeds World 4i, May 1997).

In Table 1 are listed 28 mapped and established SSLPs between Ler and Col. A number of PCR primer pairs are described herein (SEQ ID NO:42 to SEQ ID NO:97) which also yielded SSLPs between C24 and Ler (19 SSLPs) or between C24 and Col (25 SSLPs), respectively. Polymorphic SSLPs can be used as molecular markers in the analysis of homeologous recombination between genomes from these subspecies.

The PCR reactions (25 µL) were carried out over 35 cycles (15 seconds at 94°C, 30 seconds at 55°C and 30 seconds at 72°C), with 0.25 U Taq DNA polymerase and 0.6 µg genomic DNA in reaction buffer containing 2 mM MgCl<sub>2</sub>. PCR products were separated by agarose gel electrophoresis (4% ultra high resolution agarose) and visualised by ethidiumbromide staining. The results from the PCR experiments are summarised in

Table 1, which also shows the sequence of PCR primers, primer annealing temperature (Tm), PCR product length and chromosome location of SSLP (with respect to the RI map of May 1997, Weeds World 4i).

### F. Production of hybrid plants

5 C24 plants heterozygous for chlorina3A:Ac/GUS are crossed as male to emasculated wildtype Ler to produce Ler/C24(chlorina3A, GUS) hybrid seeds.

Due to the heterozygosity of the C24 parent, only 50 % of hybrid plants have inherited the chlorina3A:Ac/GUS locus. The remaining 50% of hybrid plants are wildtype with respect to chlorina3A:Ac/GUS. Since the mutant locus is linked to a kanamycin resistance gene (contained on the same T-DNA as GUS) mutant plants can be pre-selected by germinating hybrid seeds on germination medium containing 50 mg/L kanamycin.

Ler plants homozygous for the five chromosome markers are male sterile (ms1-1) and are crossed without emasculation to wildtype C24 to produce Ler(an-1, py-1, gl1-1, cer2-1, ms1-1)/C24 hybrid seeds.

Other Ler plants, which are male fertile, are crossed after emasculation of the female parent to produce Ler/C24 hybrid seeds.

G. Introduction of MSH3 and MSH6/3 antisense genes into Arabidopsis and analysis of meiotic homeologous recombination

## (i) Transformation of hybrid plants and analysis of homeologous meiotic recombination

The plant transformation vectors comprising the antisense genes described in (B) and (C) above are introduced into Agrobacterium tumefaciens strain AGL1 (Lazo et al., 1991, Bio/Technology 9, 963-967) by electroporation. Recombinant Agrobacterium clones are selected on LB medium containing 50 mg/L rifampicin and 100 mg/L carbenicillin. Selected clones are used to infect roots of Arabidopsis hybrid plants (described in (F) above) using the root transformation protocol of Valvekens et al. (1988, PNAS 85, 5536-5540) except that shoot and root inducing media contain hygromycin (10 mg/L) instead of kanamycin.

Plants regenerated from roots of hybrid plants are genetic clones of root donating plants and therefore are again genetic hybrids of two Arabidopsis subspecies described in 30 (F). However, in contrast to the root donating plants, the regenerated hybrid plants also contain the introduced transgene and the co-introduced hygromycin resistance gene with the latter allowing these plants to grow on hygromycin containing culture medium.

Hygromycin resistant plants are then allowed to enter the reproductive phase and to produce gametes by meiotic divisions of microspore and megaspore mothercells. At meiosis, the *DMC*1 promoter is activated and can direct the expression of antisense genes described in (B) and (C) above, leading to decreased *MSH*6 and/or *MSH*3 gene

expression. This in turn depletes the gamete mothercells of MSH6 and/or MSH3 protein, thus causing alteration of MMR during meiotic divisions and increasing the recombination frequency between homeologous chromosomes.

Transgenic plants are then allowed to self-fertilise and to produce seeds. These seeds (F2 seeds with respect to hybrid production), and the plants derived therefrom, carry the homeologous recombination events which can be identified by using the visual and molecular recombination markers described in (E) above.

In case of homeologous recombination between chromosomes of *Ler* and C24(chlorina3A:Ac, GUS), the analysis concentrates on chromosome 2 by selecting plants showing the visual phenotypic marker chlorina. This marker thus serves as a reference point as it indicates that respective chromosomes 2 originate from C24. Other markers, such as GUS or molecular markers, on chromosome 2 may then be used to identify chromosomal regions which are derived from the *Ler* chromosome as a result of homeologous recombination. F2 plants of control transformants not expressing the antisense gene(s) can be analysed in parallel and the results can be used for comparison to homeologous recombination results obtained in antisense plants.

# (ii) Transformation of C24 wildtype, hybrid plant production and analysis of homeologous meiotic recombination

Introduction of MMR altering genes into wildtype C24 is done using the root transformation protocol as described in G(i) for transformation of hybrid plants. Transformed plants are selected by resistance to either 10 mg/L hygromycin (in case of transformation with T-DNA's derived from pNOS-Hyg-SCV) or to 50 mg/L kanamycin (in case of transformation with T-DNA's derived from pBIN19).

Transgenic plants are then allowed to self-fertilise and to produce seeds (T1 seeds).

Segregation of the antibiotic resistance gene in the T1 population then indicates the number of transgene loci. Lines with a single transgene locus (indicated by a 3:1 ratio of resistant:sensitive plants) are selected and are bred to homozygosity. This is done by collecting selfed seeds (T2) from T1 plants and subsequent testing of at least four independent T2 seed populations for segregation of the antibiotic resistance gene. T2 populations which do not segregate the antibiotic resistance gene are assumed to be homozygous for both the resistance gene and the linked MMR altering gene.

C24 plants homozygous for the MMR altering gene are then crossed to Ler lines homozygous for recessive visual markers (see E(ii)) to produce C24/Ler hybrid plants as described in (F). These F1 hybrids are then allowed to enter the reproductive phase and to produce gametes by meiotic division of microspore and megaspore mothercells. At meiosis, the DMC 1 promoter is activated and can direct the expression of antisense or sense genes described in (B), (C) and (D) above, leading to decreased MSH6 and/or MSH3 gene expression. This in turn depletes the gamete mothercells of MSH6 and/or MSH3

protein, thus causing alteration of MMR during meiotic divisions and increasing the recombination frequency between the homeologous chromosomes of C24 and Ler. Recombination events are then scored in the F2 generation.

For recombination analysis, the hybrid plants are allowed to self-fertilise and to produce F2 seeds. F2 plants are then preselected for a first visual marker. Since this marker is recessive, its visual presence indicates homozygosity for Ler DNA at the relevant locus. Those F2 plants which show this first visual marker are then analysed for the presence or absence of a second visual marker which in the Ler parent is closely linked to the first marker. Absence of the second visual marker indicates recombination between the relevant C24 and Ler chromosomes between the first and second marker. The frequency of recombination in transgenic hybrids is compared to the recombination frequency in control hybrids not expressing the MMR altering gene.

Examples of recombination analysis are the following.

The Ler line NW22(dis1, ga4, th1) is used for crosses to transformed C24.

F2 plants are preselected first for thiamine requirement (th1) and then are further analysed for re-appearance of wildtype height (loss of ga4) and/or re-appearance of normal trichomes (loss of dis1) as a result of recombination.

The Ler line NW10(tz-201, cer3) is used for crosses to transformedC24.

F2 plants are then preselected first for thiazole requirement (tz) and then are further analysed for re-appearance of normal, i.e. non-shiny stems (loss of cer3) as a result of recombination.

The Ler line NW134 (ttg, ga3) is used for crosses to transformed C24. F2 plants are first preselected for dwarfish appearance (ga3) and are then analysed for re-appearance of trichomes (loss of ttg) as a result of recombination.

Ler plants homozygous for abi3-1 and gl1-1 are used for crosses to transformed C24. F2 plants are first preselected for their ability to germinate in the presence of abscisic acid and are then analysed for re-appearance of trichomes on the leaves (loss of gl1-1) as a result of recombination.

In the case of homeologous recombination between transformed C24 and the Ler line NW1 (an-1, py-1, gll-1, cer2-1, msl-1), recombination analysis is similar the one described above, except that the second marker is not a visual marker but has to be a molecular marker. This is because the Ler parent carries only one visual marker per chromosome.

	T	ABLE 1:	SSLP Markers in Arabidopsis thaliana Subspecies	haliana S	ubspecies		
Сһготоѕоте	RI Map Position	PCR Primer Pair	Primer Sequence	Tm l°C!	length/COL	length/LER	length/C24
	2.3	ATEAT! F	GCCACTGCGTGAATGATATG	57.8	172	162	162
		ATEAT! R	CGAACAGCCAACATTAATTCCC	58.2			
_	9.3	NGA63 F	AACCAAGGCACAGAAGCG	57.3	111	68	120
		NGA63 R	ACCCAAGTGATCGCCACC	59.6			
	40.1	NGA248 F	TACCGAACCAAAACACAAAGG	56.1	143	129	no amplific.
		NGA248 R	TCTGTATCTCGGTGAATTCTCC	58.2			
	81.2	NGA128 F	GGTCTGTTGATGTCGTAAGTCG	60.1	180	061	no amplific.
		NGA128 R	ATCTTGAAACCTTTAGGGAGGG	58.2			
	81.2	NGA280 F	CTGATCTCACGGACAATAGTGC	1.09	105	85	85
		NGA280 R	GGCTCCATAAAAAGTGCACC	57.8			
_	111.4	NGAIII F	CTCCAGTTGGAAGCTAAAGGG	93	128	162	170
		NGAIII R	TGTTTTTAGGACAAATGGCG	70			
=	ca. 7.5	NGA168 F	CCTICACATCCAAAACCCAC	57.8	213	217	208
		NGA 168 R	GCACATACCCACAACCAGAA	57.8			

3 11							
				0 23	101	001	36
	ca. 48	NGA1126L	CGCTACGCTTTTCGGTAAAG	87.8	161	3	8
		NGA1126R	GCACAGTCCAAGTCACAACC	59.9			
9	62.2	NGA361L	AAAGAGATGAGAATTTGGAC	51.7	114	120	114
		NGA361R	ACATATCAATATATTAAAGTAGC	49.5			
11	73	NGA168 F	TCGTCTACTGCACTGCCG	9.69	151	135	135
		NGA168 R	GAGGACATGTATAGGAGCCTCG	6.19			
0	ca. 77	AthBIO2 L	TGACCTCCTCTTCCATGGAG	59.9	141	209	139
		AthBIO2 R	TTAACAGAAACCCAAAGCTTTC	54.5			
II	ca. 83	AthUBIQUE L	AGGCAAATGTCCATTTCATTG	54.1	146	148	148
		AIhUBIQUE R	ACGACATGCCAGATTTCTCC	57.8			
=	3.4	NGA172 F	AGCTGCTTCCTTATAGCGTCC	35	162	136	140
		NGA172 R	CATCCGAATGCCATTGTTC	55.4			
=	12.8	NGA126 F	GAAAAACGCTACTTTCGTGG	56.1	119	147	no amplific.
		NGA126 R	CAAGAGCAATATCAAGAGCAGC	58.2			
=	17.5	NGA162 F	CATGCAATTTGCATCTGAGG	55.8	107	89	no amplific.
		NGA162 R	CTCTGTCACTCTTTTCCTCTGG	60.1			

	×	NGA6 F	TGGATTTCTTCCTCTTCAC	56.1	143	123	143
		NGA6 R	ATGGAGAGCTTACACTGATC	56.1			
21	19.8	NGA12 F	AATGTTGTCCTCCCTCCTC	6.65	247	234	220
		NGA12 R	TGATGCTCTCTGAAACAAGAGC	58.2			
2	24.1	NGA8 F	GAGGCAAATCTTTATTTCGG	56.1	154	198	190
		NGA8 R	TGGCTTTCGTTTATAAACATCC	54.5			
<u>\</u>	102	NGA1107 L	GCGAAAAACAAAAAATCCA	50.2	150	140	140
		NGA1107 R	CGACGAATCGACAGAATTAGG	58			
>	11.8	NGA225 F	GAAATCCAAATCCCAGAGAGG	58	611	189	611
		NGA225 R	TCTCCCCACTAGTTTTGTGTCC	1.09			
>	16.7	NGA249 F	TACCGTCAATTTCATCGCC	55.4	125	115	115
		NGA249 R	GGATCCCTAACTGTAAAATCCC	58.2			
				· ,			
>	19.9	CA72 F	AATCCCAGTAACCAAACACACA	56.3	124	011	110
		CA72 R	CCCAGTCTAACCACGACCAC	61.9			
>	20	NGA151 F	GTTTTGGGAAGITTTGCTGG	55.8	150	120	130
		NGA151 R	CAGTCTAAAAGCGAGAGTATGATG	58.6			

V         24         NGA106 R         GTTATGGGGTTTCTAGGGCACG         60.1         157         123           V         37.8         NGA106 R         TGCCCCATTTGTTCTTCT         55.8         174         132           V         37.8         NGA139 F         AGAGCTACCAGATCCGATGG         59.9         174         132           V         50         NGA139 R         GGTTTCGTTTCACTATCCAGG         55.8         174         132           V         50         NGA76 F         GGAGAAATGTCACTCCCACC         60.1         231         >250           V         61.1         ATHSO191 L         CTCCACCAATCATGCAAATG         55.8         148         156           V         61.1         ATHSO191 R         TGATGTTGAAGGAGATGGTCA         53.8         148         156           V         61.1         ATHSO191 R         TGATGTTGAAGGAGATGGTCA         53.7         179           V         81.7         NGA129 F         TCAGGAGGAACTAAAGTGAGG         60.1         177         179           V         81.7         NGA129 R         CACACTGAAGGTCTGAGG         60.1         177         179								
NGA106 R TGCCCATTTTGTTCTTCTC	^	24	NGA106 F	GTTATGGAGTTTCFAGGGCACG	1.09	157	123	03.1
NGA139 F         AGAGCTACCAGATCCGATGG         59.9         174           NGA139 R         GGTTTCGTTTCACTATCCAGG         55.8         174           NGA16 F         GGAGAAATGTCACTCTCCACC         60.1         231           NGA76 R         AGGCATGGAGACATTTACG         57.8         148           ATHS0191 L         CTCCACCAATCATGCAAATG         55.8         148           ATHS0191 R         TGATGTTGATGGAGATGGTCA         53.7         177           NGA129 F         TCAGGAGGAACTAAAGTGAGG         60.1         177           NGA129 R         CACACTGAAGATGGTCTTGAGG         60.1         177			NGA 106 R	TGCCCCATTITGTTCTCTC	55.8			
NGA139 F         AGAGCTACCAGATCCGATGG         59.9         174           NGA139 R         GGTTTCGTTTCACTATCCAGG         55.8         174           NGA16 F         GGAGAAATGTCACTCTCCACC         60.1         231           NGA76 R         AGGCATGGAGACATTTACG         57.8         148           ATHS0191 L         CTCCACCAATCATGCAAATG         55.8         148           ATHS0191 R         TGATGTTGATGGAGATGGTCA         53.7         177           NGA129 F         TCAGGAGGAACTAAAGTGAGG         60.1         177           NGA129 R         CACACTGAAGATGGTCTTGAGG         60.1         177								
NGA139 F         AGAGCTACCAGATCCGATGG         59.9         174           NGA139 R         GGTTTCGTTTCACTATCCAGG         55.8         174           NGA16 F         GGAGAAATGTCACTCTCCACC         60.1         231           NGA76 R         AGGCATGGGAGACATTACG         57.8         148           ATHS0191 L         CTCCACCAATCATGCAAATG         55.8         148           ATHS0191 R         TGATGTTGATGGAGATGGTCA         53.7         177           NGA129 F         TCAGGAGGAACTAAAGTGAGG         60.1         177           NGA129 R         CACACTGAAGATGGTCTTGAGG         60.1         177								
NGA139 R         GGTTTCGTTTCACTATCCAGG         55.8           NGA76 F         GGAGAAATGTCACTCTCCACC         60.1         231           NGA76 R         AGGCATGGAACATTTACG         57.8         148           ATHS0191 L         CTCCACCAATCATGCAAATG         55.8         148           ATHS0191 R         TGATGTTGATGGAGATGGTCA         53.7         177           NGA129 F         TCAGGAGGAACTAAAGTGAGG         60.1         177           NGA129 R         CACACTGAAGATGGTCTTGAGG         60.1         177	^	37 R	NGA139 F	AGAGCTACCAGATCCGATGG	59.9	174	132	132
NGA76 F GGAGAAATGTCACTCTCCACC   60.1   231     NGA76 R AGGCATGGGAGACATTTACG   57.8   148			NGA139 R	GGTTTCGTTTCACTATCCAGG	55.8			
NGA76 F         GGAGAAATGTCACTCTCCACC         60.1         231           NGA76 R         AGGCATGGGAGACATTTACG         57.8           .1         ATHSO191 L         CTCCACCAATCATGCAAATG         55.8         148           .7         ATHSO191 R         TGATGTTGATGGAGATGGTCA         53.7         177           .7         NGA129 F         TCAGGAGGAACTAAAGTGAGG         60.1         177           .7         NGA129 R         CACACTGAAGATGGTCTTGAGG         60.1         177								
NGA76 F   GGAGAAATGTCACTCTCCACC   60.1   231     NGA76 R   AGGCATGGGAGACATTTACG   57.8								
NGA76 R         AGGCATGGGAGACATTTACG         57.8           ATHS0191 L         CTCCACCAATCATGCAAATG         55.8         148           ATHS0191 R         TGATGTTGATGGAGATGGTCA         53.7         177           NGA129 F         TCAGGAGGAACTAAAGTGAGG         60.1         177           NGA129 R         CACACTGAAGATGGTCTTGAGG         60.1         177	>	50	NGA76 F	GGAGAAATGTCACTCTCCACC	60.1	231	> 250	200
ATHSO191 L         CTCCACCAATCATGCAAATG         55.8         148           ATHSO191 R         TGATGTTGATGGAGATGGTCA         53.7           NGA129 F         TCAGGAGGAACTAAAGTGAGGG         60.1         177           NGA129 R         CACACTGAAGATGGTCTTGAGG         60.1         177			NGA76 R	AGGCATGGGAGACATTTACG	57.8			
ATHSO191 L         CTCCACCAATCATGCAAATG         55.8         148           ATHSO191 R         TGATGTTGATGGAGATGGTCA         53.7           NGA129 F         TCAGGAGGAACTAAAGTGAGGG         60.1         177           NGA129 R         CACACTGAAGATGGTCTTGAGG         60.1         177								0
ATHSOI9I L         CTCCACCAATCATGCAAATG         55.8         148           ATHSOI9I R         TGATGTTGATGGAGATGGTCA         53.7         148           NGA129 F         TCAGGAGGAACTAAAGTGAGG         60.1         177           NGA129 R         CACACTGAAGATGGTCTTGAGG         60.1         177								
ATHSO191 R TGATGTTGATGGAGATGGTCA 53.7  NGA129 F TCAGGAGGAACTAAAGTGAGGG 60.1  NGA129 R CACACTGAAGATGGTCTTGAGG 60.1	>	61.1	ATHSO191 L	CTCCACCAATCATGCAAATG	55.8	148	156	146
NGA129 F TCAGGAGGAACTAAAGTGAGGG 60.1 177 NGA129 R CACACTGAAGATGGTCTTGAGG 60.1			ATHSO191 R	TGATGTTGATGGAGATGGTCA	53.7			
NGA129 F TCAGGAGGAACTAAAGTGAGGG 60.1 177 NGA129 R CACACTGAAGATGGTCTTGAGG 60.1								
NGA129 F TCAGGAGGAACTAAAGTGAGGG 60.1 177 NGA129 R CACACTGAAGATGGTCTTGAGG 60.1								
CACACTGAAGATGGTCTTGAGG	>	81.7	NGA 129 F	TCAGGAGGAACTAAAGTGAGGG	60.1	171	179	172
			NGA129 R	CACACTGAAGATGGTCTTGAGG	60.1			

#### **CLAIMS**

1. An isolated and purified DNA molecule comprising a polynucleotide sequence encoding a polypeptide functionally involved in the DNA mismatch repair system of a plant.

- 2. A DNA molecule according to claim 1 wherein said polypeptide is homologous to a mismatch repair polypeptide of a yeast or of a human.
- 3. A DNA molecule according to claim 1 wherein said polypeptide is homologous to AtMSH3 (SEQ ID NO: 19) or to AtMSH6 (SEQ ID NO: 31).
- 4. An isolated and purified polypeptide functionally involved in the DNA 10 mismatch repair system of a plant.
  - 5. A polypeptide according to claim 4 which is homologous to a mismatch repair polypeptide of a yeast or of a human.
- 6. An isolated and purified polypeptide selected from the group consisting of a polypeptide encoded by the gene AtMSH3 (SEQ ID NO: 18), a polypeptide encoded by the gene AtMSH6 (SEQ ID NO:30), polypeptides homologous to a polypeptide encoded by the gene AtMSH3 (SEQ ID NO: 18) and polypeptides homologous to a polypeptide encoded by the gene AtMSH6 (SEQ ID NO:30).
- 7. An isolated and purified DNA molecule comprising a polynucleotide sequence selected from the group consisting of (i) a sequence encoding a polynucleotide which is 20 capable of interfering with the expression of a plant polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human and thereby disabling said plant polynucleotide sequence; and (ii) a sequence encoding a polypeptide capable of disrupting the DNA mismatch repair system of a plant.
- 8. A DNA molecule according to claim 7 comprising a polynucleotide sequence encoding a polynucleotide capable of interfering with the expression of a plant polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human and thereby disabling said plant polynucleotide sequence.
- 9. A DNA molecule according to claim 8 wherein said polynucleotide is capable 30 of interfering with the expression of a plant polynucleotide sequence is a sense polynucleotide, an antisense polynucleotide or a ribozyme.
  - 10. A DNA molecule according to claim 7 comprising a polynucleotide sequence encoding a polypeptide capable of disrupting the DNA mismatch repair system of a plant.

- 11. A DNA molecule according to claim 10 wherein said polypeptide is homologous to AtMSH3 (SEQ ID NO: 19) or to AtMSH6 (SEQ ID NO: 31).
- 12. A DNA molecule according to claim 10 further comprising a regulation element capable of causing overexpression of said polypeptide in a cell of said plant.
  - 13. A chimeric gene comprising:
- a DNA sequence selected from the group consisting of (i) a sequence encoding a polynucleotide capable of interfering with the expression of a plant polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human and thereby disabling said plant polynucleotide sequence, and (ii) a sequence encoding a polypeptide capable of disrupting the DNA mismatch repair system of a plant; and

at least one regulation element capable of functioning in a plant cell.


- 14. A chimeric gene according to claim 13 wherein said regulation element is selected from constitutive, inducible, tissue type specific and cell type specific promoters.
- 15. A chimeric gene according to claim 13 comprising a DNA sequence encoding a polypeptide capable of disrupting the DNA mismatch repair system of a plant, wherein said regulation element is capable of causing overexpression of said polypeptide in a cell of said plant.
- 16. A chimeric gene according to claim 13 wherein said regulation element is selected from the group consisting of 35S, NOS, PR1a, AoPR1 and DMC1.
  - 17. A plasmid or vector comprising a chimeric gene according to any one of claims 13-16.
  - 18. A plant cell stably transformed, transfected or electroporated with a plasmid or vector according to claim 17.
- 19. A plant comprising a cell according to claim 18.
  - 20. A plant according to claim 19 selected from plants of the families Brassicaceae, Poaceae, Solanaceae, Asteraceae, Malvaceae, Fabaceae, Linaceae, Canabinaceae, Dauaceae and Cucurbitaceae.
- 21. A process for at least partially inactivating a DNA mismatch repair system of a plant cell, comprising transforming or transfecting said plant cell with a DNA molecule according to any one of claims 1-3 or 7-12 and causing said DNA sequence to express said polynucleotide or said polypeptide.
  - 22. A process for at least partially inactivating a DNA mismatch repair system of a plant cell, comprising transforming or transfecting said plant cell with a chimeric gene

according to any one of claims 13-16 and causing said DNA sequence to express said polynucleotide or said polypeptide.

- 23. A process for at least partially inactivating a DNA mismatch repair system of a plant cell, comprising transforming or transfecting said plant cell with a plasmid or vector according to claim 17 and causing said DNA sequence to express said polynucleotide or said polypeptide.
- 24. A process for increasing genetic variation in a plant comprising obtaining a hybrid plant from a first plant and a second plant, or cells thereof, said first and second plants being genetically different; altering the mismatch repair system in said hybrid plant; permitting said hybrid plant to self-fertilise and produce offspring plants; and screening said offspring plants for plants in which homeologous recombination has occurred.
- 25. A process according to claim 24 wherein a first gene is incapacitated in said first plant, a second gene is incapacitated in said second plant, and said first and second genes are incapacitated in said hybrid plant thereby altering the mismatch repair system of said hybrid plant.
  - 25. A process according to claim 25 wherein said incapacitation of the mismatch repair system of said hybrid plant is reversible.
- 26. A process according to claim 24 wherein a new genetic linkage of a desired characteristic trait or of a gene which contributes to a desired characteristic trait is 20 observable in at least one of said offspring plants.
- 27. A process for obtaining a plant having a desired characteristic, comprising altering the mismatch repair system in a plant, cell or plurality of cells of a plant which does not have said desired characteristic, permitting mutations to persist in said cells to produce mutated plant cells, deriving plants from said mutated plant cells, and screening said plants for a plant having said desired characteristic.
- 28. A process according to claim 27 wherein said step of altering the mismatch repair system comprises introducing into said hybrid plant, plant, cell or cells a chimeric gene according to claim 13 and permitting the chimeric gene to express a polynucleotide which is capable of interfering with the expression of a plant polynucleotide sequence in a mismatch repair gene of the hybrid plant, plant, cell or cells, or a polypeptide capable of disrupting the DNA mismatch repair system of the hybrid plant, cell or cells.
  - 29. A process according to claim 28 comprising inactivating an MSH3 gene and/or an MSH6 gene of said plant.
- 30. A process according to claim 28 comprising inactivating an MSH3 gene and an MSH6 gene of said plant.

- 31. A process according to claim 27 comprising at least partially inactivating the mismatch repair system of said plant in a predetermined cell type or in a predetermined tissue of said plant.
- 32. A process according to claim 31 further comprising restoring mismatch repair 5 in said cell type or said tissue.
  - 33. An oligonucleotide capable of hybridising at 45°C under standard PCR conditions to a DNA molecule according to claim 1 with the proviso that said oligonucleotide is other than SEQ ID NO:1 or SEQ ID NO:2.
- 34. An oligonucleotide capable of hybridising at 45°C under standard PCR to conditions to the DNA of SEQ ID NO: 18 with the proviso that said oligonucleotide is other than SEQ ID NO:1 or SEQ ID NO:2.
  - 35. An oligonucleotide capable of hybridising at 45°C under standard PCR conditions to the DNA of SEQ ID NO:30 with the proviso that said oligonucleotide is other than SEQ ID NO:1 or SEQ ID NO:2.

Figure 1






Figure 2

Comments/References: 52= 3' side of S5 (AtMSH3) 1244bp in pUC18/Sma1




Figure 3

Comments/References: 13 = 3' side of S5 (AtMSH3) 2104bp in pUC18/Sma1

Figure 4

08	144 15	204 35	264 55	324 75	384 95	115	504 135	564 155	624 175	68 <b>4</b> 195	744	804 235	864 255	924 275
AGCGCGCGAAAATTGGCAACCCAAGTTCGCCATAGCCACGACCACGACCTTCCATTTCTCTTAAACGGAGGA	CCC P	CCG P	CTC	GAT D	GTT V	AAG K	GGA G	TTC	GCA	GCA A	GAA E	AAT N	gaa E	GTT V
ACGC	GCT A	CCA P	CAC H	CCC	TAT Y	CTA L	TTC	N N	AAT	GGT G	CTT	AGT S	ATT	GTT V
TTA	TTC F	CCG	GAC D	GTA V	GAA E	GAG E	TTC F	CAC H	GTG V	CAT H	ACG	CAG	GGT G	GAA
TCTC	TTC F	ACA T	TCC S	CCA P	GAG E	GTG V	AGA R	GAT D	CTG L	TCC	9 8 8	TCA	TGT	GGT G
CAT	CGT R	TCA S	CTC L	AAC	CCG P	GTG V	TAC Y	ATG M	AGA R	AAG K	AAA K	GGT	၁၅၅	ACA T
CTTC	TCT	TCA S	CTT L	CAA	TCG S	CAA	AGG R	CAT H	AGA R	AT'I I	ACC	TTT F	TTA L	TCG S
ACGAC	ATT	GAA E	AAG K	ACT	CCC	CAG Q	TAC Y	GCT A		GCC	TAT Y	GGT G	ACA T	ATT I
3ACC/	ACG T	9 8 8 8	CGT R	CAC	GAA E	GAA	GGT G	TAC Y	CAT H	GCA A	TTG L	GAA E	GAG E	GAA E
CCACC	CAG	GTA V	AAG K	CCT P	CTG L	TTG L	GTT V		TTC F	ACT	GCG A	GAA	TCG	GTT V
ATAGO	CAG Q	CCG P	TCC S	TCT S	TTT F		GAA E	GGT	AAT	GAA E	TCG S	GGT G	A.A.G K	၁၅၅
מפכמ	AAG K	AAT N	CCT	CTT L	AGA R	ACA	GTG V	TTG L	TTG L	ACT T	CTG L	GGT G	GTT V	GTT V
4GTT(	CAA O	CCG	TCT	A AA	CAG O	TAC Y	ATG M	GTG V	CGA R	CAG Q	GCA G	TGT C	AGA R	GTT V
CCCA	A'A'G	GAA E	TTC	CCT	CTC	AAA K	TTG L	CGC R	TTT F	A.A.G K	CGG R	GGT G	GAG E	GGT G
SCAA	၁၅၅	CAC H	TCC	AAG K	TTT F	AGG R	GTT V	GCA A	ACA T	GTG V	TTC F	GGT G	GAT D	GTC V
ATTG	ATG M	ACT T	GTA V	AAA K	AGA R	TCG S	GTG V	GCA A	CCA P	GTA V	TTT F	AGT S	GTG V	AGA R
SAAA	ATAAAGCAATT	CCG	ACT	CCC	CAA	TCA S	GAT D	ATC I	GTG V	GGT G	CCT P	ATA I	GTT V	GTT V
2525	AAGC	TCC S	gcc •	TCA S	CAC H	TCA S	CCA P	GAG E	AGT S	ATT I	၁၅၅	GAT D	TGT C	GAT D
	AATA	X AA	TCC S	606 <b>A</b>	TTA L	ACG T	TAC Y	<b>9</b> 00	GCG A	A.A.G K	ACC	GAG E	GTT V	TTT F
cCTAAGA	GATTACG	CCC P	ATA I	200 <b>4</b>	AAT N	GAA E	AAG K	GAC D	ACG T	TAC Y	CGG R	GCT A	$_{\rm L}^{\rm TTG}$	AGT S
CCT	GAT	AAA	AAG K	. <b>₹</b>	CCC P	CCC	AGC	GAA	ATG M	GGA G	AAC	GCG A	T'IC F	ATG
ਦਜ	91	145	205	265	325 76	385 96	445	505 136	565 156	625 176	665 196	745	805 236	865 256

984 295	1044 315	1104 335	1164 355	1224 375	1284 395	1344 415	1404 435	1464 455	1524 475	1584 495	1644 515	1704 535	1764 555	1824 575
TTG L	GTG V	AGC	TTA L	. GTT	CAT	TCA	N	GTA V	ATA	TCC S	F	GGA	GCT A	AGG R
AGC S	TTG L	TTC F	AAC N	ACA T	<b>TG</b> C C	TTG L	A A A	ACA	TTG L	TCT	GAG E	CGT R	GAA E	ATG M
TTG L	TTT F	TGT	GGT G	TTG L	TTT F	TCT	GTG V	CTT L	AAT N	AGT S	CCT P	CAN	ATG M	GAA E
ATT	A.A.G. K	GAT D	GCA	76C C	ACG T	CGC R	GTT V	ACA	AGA R	CAT H	TCA S	ATT I	GTT	rcr s
GTG V	GAG E	CTG L	AGC S	TCT	CTA L	TTT F	GAG E	CAC	GAT D	TCT	GTA V	GAT D	GCA	GAC D
GCT A	ACT	TCA S	ATC I	ATG M	GCC A	TCA S	$_{ m L}^{ m TTG}$	AAT N	7.GC C			TCT	ATT	GGC ATA AAG CAA G I K Q 1 (Continued)
GAG E	CAA	GCC A	AAA K	GGA G	CTC L	GCC A	CAG Q	ATG M	CTA L	ATG	GCA A	TCA S	TTC	AAG K inu
TTA L	CAA	CGT R	GAA E	A.A.A K	000 <b>4</b>	000 0	S C	AAT N	CCT P	TGC C	AGA R	AGA R	GAG E	ATA I
GGA G	TCA S	GAA E	TGT	GAA E	CA A	CAA	CTG L	CAT H	CAT H	GCT A	GAG E	TCT S	ACA T	<b>၁)</b> ၁၂၅၅
AGT S	CTT L	GTG V	TTA L	GCT A	GTT V	TAC Y	ACT	TTC F	ACT T	TCT S	TCT	ATG M	000 <b>4</b>	
AGA R	CCT P	CGA R	TCA S	GCT	ACT T	CTT L	AAT	TŤA L	GTG V	ATT I	GGT G	GCT	AAA K	cgg crr R L
ATG M	CAG Q	GTT V	ATT I	GAG E	CTG L	ATC I	GCC A	TCC	TGG W	GAG E	GAA E	ACA T	GCT A	CAG Q
TTC F	၁၅၅	AAC N	GTT V	CTG L	CAT H	AGG R	TCA S	၁၅၅	CAC	TCT	GAA E	$_{\rm L}^{\rm TIG}$	ACT	ATT I
AAT N	CTT L	TCA S	GAG E	AAG K	CCA P	GAA E	CTC L	TCT	AGA R	GTT V	GTT V	GTC V	CGG R	CAA O
GAT D	${f TTG}$	ACC T	GAT D	ATG	ATG M	TTT F	ACT T	GAA E	CTT L	GCT A	${ m TTG}$	TCA S	CAT H	A.A.G
AAT	CTG L	CCT	GTA V	GAA E	AAC	GGA G	ATG M	TCG S	CTT L	GAT D	GAG E	TCC S	TTT F	9 9
TTC	GAG E	GGA G	GCA	AAA K	ATG M	ttt F	GAG E	GGA G	AGG R	CTT	AGT S	CTC L	ATC I	6CG ▶
GAG E	GCT A	GCT A	AAT	GAT D	ATT I	CAG O	ACA T	GAT D	TCC S	CGG R	AGC S	GTG V	AGA R	CTT L
GAA E	CCA P	CAT M	GGT	GAT D	ACA T	A AAA	AAC N	TCA S	GCT G	GCT A	CTC L	CTC L	ACA T	TTA L
TAT	TCA S	GCA	AAT	GAA	CAT H	CTC L	AGT S	AAT N	TAT Y	TCT	CAG Q	TAT Y	ATA I	ATT I
925 276	985 296	1045 316	1105 336	1165 356	1225 376	1285 396	1345 416	1405 436	1465 456	1525 476	1585 496	1645 516	1705 536	1765 556


1884 595	1944 615	2004 635	2064 655	2124 675	2184 695	2244 715	2304 735	2364 755	2424 775	2484 795	2544 815	2604 835	2664 855	2724 875
-	<u>د</u>	£.		C)	(J)	æ	£ ·	U	E	Æ	U	æ	£+	4
TC.	GCT A	GCT A	TTT E	TTG L	A.A.G	GCA	AGT	CAC	TGT	CAA	ATC	ATA I	GGT G	GAA E
ATT I	gcg A	CTT L	TCA	CAT H	ACC T	CTA L	TTC F	TTG L	GAC	TTA L	ATT	TCC	GAT D	CTA L
GTT V	GAA E	GAG	GCT	ACA	AGC	GCT A	AGT	TGT C	GAT D	ATA I	C.P.A	ATT I	CTT L	TTT F
TCT	AAG K	CCT	ATA I	ATC I	AAT	CTA	AAG K	GAC D	GTG V	ACT	16C C	TTA L	GTG V	ACC T
ATT	AAT N	TTT F	TCG S	999	GTA V	GAG	CTC	CTG L	TTT F	GAG	TAT Y	GCT A	CAC	AGT S
TTG L	CTA L	CAA	TCC S	TCG S	A AAA	GAT D	TTC F	GCA A	GAG E	CTG L	GAA E	GTT V	ĊTG L	AGA R Bd)
AAA K	GCC A	GAC D	GAT	GTG V	GTG V	T16	AGT S	GCT A	CCC	GTA V		CAA O	A'AG K	G CAT GGC AG H G R (Continued)
AGA R	TCT S	AGC S	CTG L	CAA	TGG ¥		GAT D	CTT L	CGT R		GAA E	CGT	9 8	CAT H
TTG L	CTC	TCC	AAG K	CTT L	A N	GCT.	TGG W	GCT A	GTC V	CAT H	GCA A	ATC I	TTC	ပ်ံဝ
CTT L	CTT L	AC'F T	GAA E	TTT F	ATG H	GTA V	TCG S	CAR O	TAT Y	CGT R	CAT H	TAT Y	TCA S	ATC I
ACT T	A X	ATC I	AGG R	GAA E	CCT	ATA I	GCT	GTT V	AAC N	GGT	TTG L	76C C	GCG A	AGT ATC S I
TCT	GGA G	CTA	ATC I	TTG L	GTC V	GAA E	CGA R	GCC A	A.A.G.	TCT	ATT I	AGC S	CCA P	GAC D
CGA R	GCC	AŢA I	GTC V	AAT N	AAG K	CCA P	AAC	GC'F A	AAC	CAG Q	ACA T	AAG K	GTA V	TCAS
GTG V	A A T	GAC	TTA I.	CGA R	TCC	င် (၁	GTG V	AAG K	AGA R	ATA I	GAC D	GGA G	TTT F	GCT A
ACT	GAC D	CTC L	GTT V	ATT I	GAT D	CAT H	ATT I	TTT F	TCT S	AAC N	AAT N	GGA G	TCC S	ggt G
GCA	GTT V	TTG L	GCA	GCT A	GTT V	TAT Y	GCC	GAT D	CTA L	ATA I	CCA P	ATG M	GGT	ATG
TCT S	GTG V	GAC D	CAA	CTC L	CCC P	CGA R	$_{\rm L}^{\rm CTT}$	ACA T	ACT T	GAG E	GTC V	AAC	GTT V	000 8
CAA O	GTT V	GGT G	CGC R	A.A.G K	CTG L	ATT I	CAT H	TAC Y	TCA S	GTT V	TTC F	CCT	CAG Q	ACT
ATG M	CCT	CGA R	GCT	A.A.G K	GAG E	ACT T	GAA	TAC Y	CTT L	CCA P	AAC	GGA G	GCT A	TTC
AGT S	TCC	GTT V	GAA E	CGC	ATA I	A.A.G K	ACT T	AGA R	TCC	GAA E	GAT D	ACC	ATG M	GTT V
1825 576	1885 596	1945 616	2005 63 <b>6</b>	2065 656	2125 676	2185 696	2245 716	2305	2365 756	2425	2485 796	2545 816	2605 836	2665 856

2784 895	2844 915	2904 935	2964 955	3024 975	3084 995	3144 1015	3204 1035	32 <b>64</b> 1055	3324 1075	3397 1082	3458 5	3522 16
TTA L	TTA L	ATA I	ACA T	GTG V	CCA P	AGA R	GAA E	GAC D	CTA L	909	ပ္	
										GTT	P CCG	
ATA	ACA T	GAA E	CTG L	CTT L	CCT	GCA	GAA	GAG E	AGA R	TTA	AAT	AA
GTT V	GCA A	CCT P	TAT Y	AAG K	ATA I	CGT R	GCA	GAA	ATC	TCTG	TCT	AAA
CTT	TAT Y	TAC Y	TCG	TAT Y	cAG 0	GTA V	၁၅၅	TCT S	AAA K	ATCA	GAT	AAA
TCG S	8 8 8	CAT H	GTC V	CTA L	GCC	GAG E	AGA R	CTC L	၁၅၅	CTTG	ATG	AAAA
CGT R	ATT I	ACG T	CAT H	TAC Y	CTT L	GCT	CCG P	GCT A	GCT A	TGA TTTAATCTTAACATTATAGCAACTGCAAGGTCTTGATCATCTGTTAGTTGCG	TAG AGAG ATG	TAA AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
TCT	OCC P	GTC V	TAC Y	ACC	CAG Q	gaa E	gaa E	T'IT F	ATT 1	TGCA	AG A	GCA TAA AAAAAAAAAAAAA A Figure 4 (Continued)
TCT S	GTA V	TTT F	ACA T	GTG V	GCT	TTG L	GAA E	ξ×	AAG K	CAAC	NAT T	Con1
TGT	GGT G	CTT	ე ე	GAT D	GTT V	AAA K	CAT	CTG L	TGG	ATAG	AG P	AAA A
ACC	GAC D	GTT V	GTT V	GAT D	A.A.G K	GCA A	GGA G	GAC D	GCT A	CATT	AA :	TAA
AGA R	CAC H	TTG L	TCT	CAT H	TTT F	GCT A	GAA E	GCA A	CAT H	TTAA	GA A	GCA A Figu
ATC	ACA T	TGT	GGT G	GAT D	GGT G	ATG M	CCA P	TTT F	TTA AAG L K	AATC	CA A	cca ccc gca P P A <b>Figu</b>
ATA	AGC S	AGA R	CCA P	TAT Y	TIT F	TCA S	GAA E	TTG L		TTT	TA A T	CCA P
CAC H	ACT	A.A.G K	TTC F	AGT S	AGC S	ATT I	GGA G	GAC D	TTT F	TGA.	AGT ATA ACA AGA AAA GAG S I T R K E	TCT
TCA S	၁၅၅	GAA E	GGA G	၁၅၅	AGG R	GCC	ATG M	GGT G	GAG E	TTT F	ATT A I S	FTT
GCG A	AGA R	GCA A	AAC	A A A	AGC S	CGA R	CGC R	CTA L	TTC F	TCA S	TGT A	ATC
GAA	GGA G	CTA L	AGT S	GAT D	<b>1</b> 60	CGT R	ACA	GCT A	GCA A	TGT C	ATG T	TAC
AGT	CTT	CTC L	ATC I	A.A.G K	CTT L	ATA	AAT	TCG S	A A	ACT T	TT A	CAG Q
TTA L	GAG E	CAT H	GAG E	CAG O	GGT G	TGT C	AGA R	ATT I	TGG ₩	CCA P	TACTAACTT A	TTG L
GAA E	GAT D	CAG	GCT A	TTG L	cgt R	TCA	GAG E	TCT	CCT P	A.A.A.	TAC	GTG V
2725 876	2785 896	2845 916	2905 936	29 <b>6</b> 5 956	3025 976	3085 996	3145 1016	3205 1036	3265 1056	3325 1076	3398	3459 6

Figure 5

LOTORKOMLS SVASTTOTED STRIKES	POVŽBAVEŽOVA DAV ROMŽEV ERZOVA DA C	SSVZERKISMVFTRA	OPENANTE OF A ME	YQALALTPCHLKOPG IRLVSKLYSHVKYW,	AVERIBACHUS NA	Skos-atvastedar Isboalukospija	RETLOVEGITHLIELP Mondrovovilevens	DESTRUCTOR STATES OF THE STATE	DONETRN SASDSTONS RWZLERIGANODIIM	PYHVEYL TLOKOKUSY NYHVOKY EROK: - TUK	LOADER PER PER PER PER PER PER PER PER PER P	
SANN FILLILIA LA COPTION PERFORMANCION PROPERTO CANDA CONTRA CANDA CONTRA CANDA CONTRA CONTRA CANDA CONTRA CONTRA CANDA CANDA CONTRA CANDA CAN	PHIOMPY FORNINGS FLOORLER	ab	· · · · · · · · · · · · · · · · · · ·	SKYNYKRASIDCYSMGNAVDWY MSCHTUSACNIKDDKKMKIRARKGMSCLTVETTETT.  1. TRIVOKLTOKYDLYVQALKYKRUM MOCESTETTETTETTETTETTETTETTETTETTETTETTETTE	igatempsart coopevykan sogs krostem metlen krostem vrostem verde or mede samena verses. Igai bijlov na losudi ptili Gen - Geoppal lubem spolenjaraji er povedije spolike ite sejam s	rpylvisgvitams ms sologuitgipundakatgy iavmeatilaorqiqaie, bevustruges. ippestroment policitams my colts ar dv tytromy rauges lisenpests domi ur og promy	orlesalenten propledilits-sögepelarrandavlyzärkudsstäsperen anneselovelikut. Preden enven promopen linkudes solinkių aksesyšsos kerinkių aksesus sum kauda promosiliku	tekt ihteppelvagldköllatrherivnasmos poks fork tröperaavga okaloggabos tesker kyvakelvd Trovs kyttektok et okölt yndeleinkuselot kestonk itar it en omot ogtesjaatsonvingtende	VLETIDODNY VEHDTALHABGEYCGIIIGENSCYIRGVALKGIUNGY YSSYVAR FARBAV LOGYFTRU SABDSFOR Ites-Boy hyvehdimms penckimittgerhogkeskiroval tyrao kospyparirus sustyralikalise	ir tobsrsovildeloros hudovala katoquelare no-dvotvorik priable do svotvihe sudeloros. Er rom riselalddo vogo duogiate ya en kolode die prip predesiapeir mumbolv ekom - to	SCHANNING BERKERSTEIN	
1 MBHQK	2 SDELAAASPKKOKLBPEHIONPVPDPNIHQKFLO.	4 POSENE IAARVSOITAB WOBM I PAEDĀVIV SRIEMĪKLV POKLTISKS NPUDCN N	NULEAR DISCOCCORROFOSOS NFLVCVVDRRV Forovestevlr Thudin	6 DSQCTEKTLVAHAGOTSNVRVRRASLDCVSNGN 8 Oflevakatorgeleinkukyblkullvvrlk	1 PSRILYQCASFRBLSGNTRHTFBART BOOGEVV	HENRY OF ALVERSANCE AND STATE OF AND STATE O	LISTERSTATOR 1 DESERVATOR 1	KHANIMAAATANIO 9	S CEPVETHIGSORME 1 40 A GIANNARME	O RSTELKELSKASHI 3 DSTERVEMLDILHI	14 DEDDUTY YES DVEG DCSRSP SPRYNO DAG 1 PPSC NEWN SEA BREEK DSV 13 Debs zijenn en money went of rever dane dro en my my banden se	19 APEPLKHAWKHAGKIR DKPPCSF 12
	40 ex	1 6 1	22	9 9	9.0	4 6	80 SO	9 9	77	67	<b>6</b> 6	10.5
M SE 3 A C	E SEN E E E E E E E E E E E E E E E E E	25 25 25 25 26 27	80 M	2 3 2 3 2 6	E SE	ESE3_ACESE3_AC	Z Z	MSH3_AC MSH3_SC	M SES AU	SA CHEN CHEN CHEN CHEN CHEN CHEN CHEN CHEN	ESE3_AC	K SE 3 . AC K SE 3 . AC

Figure 6






Figure 7

Comments/References: 43= 5' side of S8 (AtMSH6) 2182 bp in pUC18/Sma1




Figure 8

Comments/References: 62= 3' side of S8 (AtMSH6) 1379bp in pUC18/Sma1

80	153	213	273	333 64	393 84	453 104	513 124	573 144	633 164	693 184	753 204	813	873	933 264
သသ	<sub>U</sub>	TCC	999 999	AGA R	CCG P	AAA K	GAT D	AAA K	GAA E	GCT A	GTA V	GAA E	AGA R	TTC
TCC	c cAG	GTT V	GAA E	GTT V	A AG	GTA V	AAT	GGT G	GTA V	CGT R	CCT P	A A	AAT N	GTT V
ATCT	3 CGC	TTG	AAG R	GAG E	TTT F	TTT E	CTG L	AAT	TCA S	CCA P	GTT V	AAG K	SCC A	GAT D
3CCA	3 CAG	GGT '	GTG V	GAT D	GGA G	A.A.G K	CCG	AAT	AGA R	CGT R	AAG K	GAG E	GAT D	CCT P
າຣຣດຕ	A ATC	AAG C	AAT	GTC	TCT	CAT H	GTT V	TCC S	CTT L	ATG M	GAT D	GGA G	AGG R	CCA P
rtti	TCTCTCAGCTCAAAACATCGTTTCTCTCTCTCTCTCTCTC	ACG 1	TTT I	TCT	CCG P	ATG M	GTT V	CGT R	GAA E	999 9	GAG E	TGT	ATC	ATA I
AAA1	TCC	ACT /	CGA 3	X AA	CTG	AT'T I	GAT D	T'T'T F	GCT A	CCA P	AAG K	GTT V	CGA R	CAC H
3GCG	CAAT	GCG A	CCA (	TCG	GTC	AAT	GAA	CAA	AGA R	ACA T	TTT F	CCG P	TCT	TTA L
CAAG	CTC!	900 (	GGA (	GTT	CGT	TCC	CGA R	CCT P	9 9	GAA	ACT	GAT D	TCT	ACC T
GACG	CTCI	ACC C	GGA (	GCT (	CGT R	TTC	AGC	AT'T I	AGT	CCA P	ATG M	CAG	GAG E	AAG K
CTAC	CACT	CCC F	AGC (	TTT (	CCG	CTG L	AGG .	GT'r V	TTC	GGT G	GAA E	CTC	CTT L	AGA R
ATTT:	CTCI	AAA X	3 2 3 3 3 3 3	CGT	GTT (	TCC	GAG	GAT. D	AGT S	CCT P	GAT D	ATG M	TGG ¥	GAT D
2252	TTCI	CAA P	) 299	GTA (	AAG K	TCG	GGA	A A T	TTT F	GTT V	GAG	X AA	GAA E	TAC Y
TITC	TCGI	TTC	999	TCT (	GAG E	GCT	TCT	GCT	GCT A	GAT D	CTG L	$\mathop{\rm CTG}_{\rm L}$	TTT E	CTT L
ATCG	AACA	TTC 1	) ) ) )	GCT 7	CCG (	GAT O	TGT	A.A.G	CAT	၁၅၅	GTT V	<b>A</b> GG R	A AA	000 <b>P</b>
GAGT	TCAA	TCT T	AGC O	GAC (	CCA (	GST (	GAT '	ATG	AAC	GAT D	CGA R	A A	ACC	GAT D
TGAG	CAGO	TTG 1	GCT P	) 255 5	ACT (	3 3 3 4	CGA (	TGT C	AGA R	GTA V	AAG K	AAC N	GGA G	GAT D
၁၁၁၅	CTCT	ATT I	GCT G	AAA O	GAT	TCC (S	GAT (	CTA 1	GAA	GGA	${ m TTG}$	TCT S	GA.A E	CCT
TTGA	AATT	TCG A			ACG C	GAA 1	GAT (			ATA I	252 <b>%</b>	GAC D	AAC N	CGT R
AAAAGTTGAGCCCTGAGGAGTATCGTTTCCGCCATTTCTACGACGCAAAGGCGAAAATTTTTGGCGCCAATCTTTCCCCCC	TTTCGAAT	AGA T R S			GGA P	GCT G	GTC (	TCA 1	ACT (	GAT D	TCT	TTG	GTA V	AGA R
4		_				394 G	454 G				694		814 225	
		erf	2	2	(4)	(*)	4 4	4, [	47.5					

993 284	1053 304	1113	1173 344	1233 364	1293 384	1353 404	1413 424	1473	1533 464	1593 484	1653 504	1713 524	1773 544	1833 564
I.	ggr	GGT G	GTT V	ATA I	9 9	TCA	ATC I	GAA E	ACG T		S	GAG	TTT F	GAG E
GAC A	TTA L	GTT V	A A A	ACT	ATC 1	TGT C	TCC	AAG K	TAT Y	ACA T	GAA	GGA G	ATT I	CTT
ATG G M	GAA E	CAG Q	TAT Y	AAT	AAC N	AAG K	0 0	CCA P	AAA K	GAT D	TCT	CTT	GAT D	N
TAT A	GCG	AGA R	GGA G	GCT A	GGA G	CAA		TCT S	AGG R	999 9	TCT	GCT A	9 9	GTA V
GAA T	GAT	TGC	CGT R	GGT G	GAG E	CTA L		GTT V	CTA L	ATG	GGT G	AGT	CAT H	ATG M
AGT G	CTA	AAA K	GCT A	AGA R	AGC S		TTT F	CAG Q	GCT A	GTA V	¥ ¥	CTT	AAG K	ACG T
AAG A	GAG E	GGA G	TTA L	GCC	GCA	ATG	AGG R	ATG M	AAG K	CAA	TTT F	GCC	CTT	T GGC CAG AC G Q T (Continued)
GTT A V	TAT Y	GTG V	CTA	A X	ACA T	A A	TTG L	TTG L	S CA	CCA P	TAC Y	GTT V	GTA V	660 6
AGT G S	CTG L	GGT G	AAG K	GCA A	TCA S	ATC I	gcc B	TTA L	GCA	GTA V	GGA G	GAT D	GAT D	8 a 6
TGG A W	GAG E	AGT S	CAA	CAA	CCA P	GAG	GCT A	GCG A	gaa E	CCA P	AAC	TGT C	G A.A	AGA ATT R I Figure
rat t Y	TAT Y	ATG M	GTG V	GAC D	ACT T	A A	TGT	. cĠA	AGA R	GCT A	TCT S	GAA E	CTA L	AGA R Fig
CAA T	TTT F	ACC T	GCA A	TCT S	TTA L	ATA I	GAC	CII	TCA S	TTG L	GAA E	AAT	AAG	CTC L
AAG C	AAA K	ATG M	GAG E	ACA T	GTA V	GCT A	GTT	GCT A	CTA L	CAG Q	ATA I	CTA L	CTA L	TGT
CAA A	999 9	AAG K	GAT D	GAA E	CAG Q	$_{\rm L}^{\rm CTT}$	TTT F	GCT A	000 0	GTA V	ATA I	GGT G	AGG R	GGT G
rca c	GTG V	ŢGG ¥	ATA I	CTA L	GTT V	CTT L	GCT A	TGT C	AAA	GCG A	AAT N	GAT D	TCT	AGG R
GCA T	AAA K	GAC D	999 9	CAG	CTA L	CAT H	TTT F	TCA S	AGT S	ACG T	AGA R	GTT V	CTG L	TAC Y
rct G	TTT F	CTT	AGT	GAG E	AAG K	GTC V	GGA G	GCA A	GAC	TCT	GTT V	GCT A	CAT H	GTT V
7. E	TTC	GAG						GAT D	TAT Y		GGA G	TGT C	AAT N	C AA
8 ×	CIT		TCT						TTA L	ACA T	GCT A	AAC N	ATT	TAC Y
AG AJ	GTG CTT TTC	CAC 7	ATC	GGA 6	ATT CCA I P	CCT	ACT	AGC GAT S D	GTG	${ m TTG}$	GCT A	TGG W	CTA L	CCA TAC P Y
934 A 265	996 285	1054	1114	1174 345		1294 385			1474 445		1594 485	1654 505	1714	1774 545

1893 584	1953 604	2013 624	2073	2133	2193 684	2253 704	2313 724	2373	2433 764	2493 784	2553 804	2613 824	2673 844	2733 864
AAC	GAT	AGT S	292 4 8		r <b>GAT</b> D	r AAA K	A GCC	A ACT T	T CAC H	T GGA G	A ACA T	T GGT G	C ATT	T CGT R
GAT D	A AA	GAA E	GGA G		ATT I	TGT C	GCA	GAA E	ATT	GCT A	, AAA K	GAT	s AGC	r CTT L
CTT	CTC	TCA S	CTC L	¥ ×	GGA G	CTC L	GAA E	GCT A	GTC V	TCT	CAG Q	<b>₽</b>	၁၅၅	CTT
TAT	CCA P	A.A.C.	CTG L	999	AGT S	A A	TTC F	AAC	GAG E	CTC L	AAT	GCA A	AGT	ACT T
AAA	CAT H	GCA A	AGA R	CTG L	AGA R	TAT Y	CAA Q	GAA E	TCT S	AGT S	CAG Q	GTT	AGC	S
TAC	TGC C	ACG T	GAA E	C'IT L	1.1.C F	CTT L	TCT S	GAT D	7GG ₩	GCA A	GAT D		AGA R	AAA K
TTG L	ATC	TTC	TTA L	GCT A	9 9 9	TTG L	CTT L	ACA T	S O		ACA T		AGA R	GGA G Lnued
ACC T	TGG .	GAA	GAC D	CC:T P	X X	AGT	TTT F	GTG V	ACT T	ATC	GCT	CCA	GCT A	ATG GGC GGA P M G G F 9 (Continued)
666 <i>I</i>	AAT	GAA	CCA	TTG L	GTG V	ATG M	TTA L	GAT D	GCA	GCA A	GAA	CAT H	GAG	_
TCA (	AGG 1	GTT (	CTT (	GTG V	ATT I	ATG M	GAG E	CAA	AGA R	TTT F	TCA S	TGG W	ე <u>ე</u> ე	AAC N Ure
CCT T	TTA A	GTA (	AAA X	TCT	CAAS	AA'I'	CTA L	AAC N	G AA	TCT	GAA E	CTA L	$_{\rm L}^{\rm CTT}$	CCA P
GGT C	CTC 1	GAT (	CAC	229	ວນນ	TCA S	ე ე	CAG	ATC I	AGA R	CCC	GGA G	CTC	GGA G
GGT O	CGA C	CTT (	CTC (	TCA	TTT F	GAA E	AGC S	TAT Y	TTT F	CTG L	TTT F	CAA	ATA I	ACG T
GAT (	AAG O	CGG (R	TAT	TCA S	GCA	AAG K	A AA	AAT	CTT L	GTC V	ATT I	ATC I	GAT D	CTG L
TGT	GGT 1	AAA L	CAG	CGA R	A.A.A K	CAG Q	GGA G	CCA P	GAA E	GAT D	GTT V	AAA K	AAT N	TTA L
AGC 1	ACT (	AAT	299	GTT V	GTT V	CTA L	GTA V	TTT F	ATC I	CTA L	CCT	CTT	CCG P	TTG L
AAT A	CCA 1	ATC I	ACT	AGC	CGA R	GCT	TTA L	GAC D	$_{\rm L}^{\rm CTT}$	7 <b>GC</b>	AGG R	ATA I	GTT V	TCA S
AAC 7	€→		ATC	TCT		${ t TTG}$	ATA I	AGC. S	ATA I	AGC S	gcc <b>A</b>	CCA	CCT	066 8
TTT P	GTT P				AAA	TTG	CCT P	GAT D	ACA	ATA I	ATG M	ეეე ე	TTG L	CCT
ATA T			ATG O	ATC #	CTG /	CTG :	CTT		CTC	ACC T	AGC S	AAA K	CAN O	CAT H
1834 A		, 4° C		74			54 0 <b>5</b>	14 25	7	2434	2494	2554 805	2614 825	2674 845

2793 884	2853 904	2913 924	2973 944	3033 964	3093 984	3153	3213 1024	3273 1044	3333 1064	3393	3453 1104	3521 5	3579	3606 28
TGT CTG GCC GTT ATC TTT GCC CAA CTT GGC TGC TAC GTG TGT GAG TCT TGC 2793	CTC GTG GAT ACT ACT ACA AGG CTT GGC GCA TCT GAT AGA ATC ATG ACA L V D T I F T R L G A S D R I M T	ACC TIT ITG GTA GAA TGC ACT GAG ACA GCG TCA GTT C'IT CAG AAT GCA ACT T F L V E C T E T A S V L Q N A T	GTA ATC CTT GAC GAA CTG GGC AGA GGA ACT AGT ACT TTC GAT GGA TAC	GCA TAC TCG GTT TTT CGT CAC GTA GAG AAA GTT CAA TGT CGG ATG CTC TTT A Y S V F R H L V E K V Q C R M L F	CAT TAC CAC CCT CTC ACC AAG GAA TTC GCG TCT CAC CCA CGT GTC ACC TCG AAA H Y H P L T K E F A S H P R V T S K	GCT TGC GCA TTC AAA TCA AGA TCT GAT TAT CAA CCA CGT GGT TGT GAT CAA GAC	TTC TTG TAC CGT TTA ACC GAG GGA GCT TGT CCT GAG AGC TAC GGA CTT CAA GTG F L Y R L T E G A C P E S Y G L Q V	ATG GCT GGA ATA CCA AAC CAA GTG GTT GAA ACA GCA TCA GGT GCT GCT CAA GCC M A G I P N Q V V E T A S G A A Q A	AGA TCA ATT GGG GUA AAC TTC AAG TCA AGT CAC CTA AGA TCT GAG TTC TCA AGT R S I G E N F K S S E L R S E F S S	GAA GAC TGG CTC AAG TCA TTG GTG GGT ATT TCT CGA GTC GCC CAC AAC AAT GCC E D W L K S L V G I S R V A H N N A	GGC GAA GAT GAC TAC GAC ACT TTG TTT TGC TTA TGG CAT GAG ATC AAA TCC TCT G E D D Y D T L F C L W H E I K S S	GIT CCC AAA TAA AIG GCT AIG ACA TAA CACTAICIGAAGCTCGITAAGICIITIGCCICICI V P K * M A M I *	TT ATT CCT CTT AAA AAA TGC TTA TAT ATC AAA AAA TTG TTT CCT CGA TTA AAA I P L K K C L Y I K K L F P R L K	AAA AAA AAA AAA AAA K K K K K K Figure 9 (Continued)
GCA ACA		GAG	GAT	ŢŢ		CAC ATG H M	CTA GTG L V	GCA CTC	ATG AAG	1G			TG	AAA AAA K K
2734 GC	0 Q 0 4 R	9 72 G		74	34 65		<b>ፊ</b> ሊ		274	334 065	. 60 a	454	522	3580

Figure 10

s to be so to se t	
648GYRSSASVLPALURRYLKONVANTOGIVRGVINIALIALORR	
11 623 HEST MESTER STEED AT DESCRIPTION HEST PRESENTATION OF THE STEED S	6667 9282 8 - 9282 2 14
162 KYTLTOSTAVOLAPVPOVAODT DA AOVRAI MENORTA CAVULINEC - PRALEMAN BRANCA DE TROPERSONAL MONTOCLAI ROS ON TROCKAI ROS ON TROCKA	PLEBER - BI
ANDROLLE BERNESE AND	10 0 A 10
113 TPPENOVESVENTATELENINVENTATELENINGENEELDENINGENEELDEN DEN GERTHAAGTE OM FYNDER REGEREN DAE BELENEELDEN LES BELENEELDEN DE SELVEN DE	TB-9KEI14
AND THE TRANSPORT OF THE PROPERTY OF THE PROPE	ACRONG-91
49 Prekvervipegriperendelfinkum rvkvjaki cegerijardile	TH-9KBKHK
TO THE TAX OF THE TAX	ACEBEG-81 E886-81

•	
TTTTTTGGTTGCTAACAATAAAGGTATACGGTTTTATGTCATCAATATAA	50
CTATATATAAAAGAAATGAAAGATATATATTGTTTTTTCATTTATCAAAC	100
AAAACAACAAGACTTTTTTTTTTTTTTTTTTTTTTTTTT	150
GATAAACGACATCGTTTAATCATTTCCCAATTTTACCCCTAAGTTTAACA	200
GATAAACGACATCGTTTAATCATTTCGCAAGCACTGATTAGGAACAGCTTT	250
ACCATTCTCATATTCCTGAACTACCTGAGTCCTCTCATTGATCTGTTTCG	300
ACCATTCTCATATTCCTGAACTACCTCTCTCTCTCTCTCT	350
ACCTCACCTCTGCTTCACACGATCCATCGCCGCAGGCTCTGTTTCTTC	400
TTCCAGCTTCTCGTGTTAATCACCGGAACCGCCGTAGATTTCCCCTTTT	450
TTCCAGCTTCTTCGTGTTAATCACCGGAACCGCGACACCGTTT	500
TGTTCGAACCGGCATCGAATTTCTTAACCGTTTGAACGGTCTTTTG	550
CTCAGAGCTGCGTTAACCGCTTCGGATCGCGTAGGTCTTCCTGTTTTCTGTTTACTGCTCCTG	600
TTTTGATTTGTGGAGAACTACTGGTTCCCAGTCTTGTCTACTACTCACACCAAA	650
GGTATCTGCTCGGCATCGTCGATGAATTGAGAGAAAGGAACAACGCGAAA	700
ATTTTATTAATCTGAGTTTTGAAATTGAGAAACGATGAAGATGAAGAATG	750
TTGTTGAGAGGATTGTGATATTTATATATACGAAGATTGGTTTCTTAGAGATG	800
ATTCGATCATCTTTTCTCCATTTTCGTCTCTGGAACGTTCTTAGAGATG	€ 850
ATTGACGACGTGTCATTATCTGATTTGCAGTTAACCAATGCTTTTTTGGGT	900
TGGATTCGTGGTACACCATATTATCCGATTTGGCTCAATGGTTTTATATA	950
AATTTGGTTTTCGGTTCGGTTATGAGTTATCATTAAAATTAAGCTAACCA	1000
AAAATTTTCGTAAAATTTATTTCGGTTTCAATTCGGATCCCTTACTTCCA	1050
GAACCGAATTATTCGAAACCGGGGTTAGCCGAACCGAATACCAATGCCTG	1100
ATTGACTCGTTGGCTAGAAAGATCCAACGGTATACAATAATAGAACATAA	1150
ATCGGACGGTCATCAAAGCCTCAAAGAGTGAACAGTCAACAAAAAAAGTT	1200
GAGCCCTGAGGAGTATCGTTTCCGCCATTTCTACGACGCAAGGCGAAAAAT	1250
TTTTGGCGCCAATCTTTCCCCCCCTTTCGAATTCTCTCAGCTCAAAACATC	1300
GTTTCTCTCACTCTCTCACAATTCCAAAAAATGCAGCGCCAGAGAT	1350
CGATTTTGTCTTCCAAAAACCCACGGCGGCGACTACGAAGGGTTTG	1400
GTTTCCGGCGATGCTGCTAGCGGCGGGGGGGGGGGGGGAGACCACGATTT	1450
AATGTGAAGGAAGGGGATGCTAAAGGCGACGCTTCTGTACGTTTTGCTGT	1500
TTCGAAATCTGTCGATGAGGTTAGAGGAACGGATACTCCACCGGAGAAGG	1550
TTCCGCGTCGTGTCCTGCCGTCTGGATTTAAGCCGGCTGAATCCGCCGGT	1600
GATGCTTCGTCCCTGTTCTCCAATATTATGCATAAGTTTGTAAAAGTCGA	1650
TGATCGAGATTGTTCTGGAGAGAGGTACTAATCTTCGATTCTCTTAATTT	1700
TGTTATCTTTAGCTGGAAGAAGAAGATTCGTGTAATTTGTTGTATTCGTT	1750
GGAGAGATTCTGATTACTGCATTGGATCGTTGTTTACAAATTTTCAGGAG	1800
CCGAGAAGATGTTCCCGCTGAATGATTCATCTCTATGTATG	1850
ATGATGTTATTCCTCAATTTCGTTCCAATAATGGTAAAACTCAAGAAAGA	1900
AACCATGCTTTTAGTTTCAGTGGGAGAGCTGAACTTAGATCAGTAGAAGA	1950
TATACGAGTAGATGGCGATGTTCCTGGTCCAGAAACACCAGGGATGCGTC	2000
CACGTGCTTCTCGCTTGAAGCGAGTTCTGGAGGATGAAATGACTTTTAAG	2050
GAGGATAAGGTTCCTGTATTGGACTCTAACAAAAGGCTGAAAATGCTCCA	2100
GGATCCGGTTTGTGGAGAGAAGAAGAAGTAAACGAAGGAACCAAATTTG	2150
AATGGCTTGAGTCTTCTCGAATCAGGGATGCCAATAGAAGACGTCCTGAT	2200
GATCCCCTTTACGATAGAAAGACCTTACACATACCACCTGATGTTTTCAA	2200
Figure 11	

Figure 11

GAAAATGTCTGCATCACAAAAGCAATATTGGAGTGTTAAGAGTGAATATA		2250
TGGACATTGTGCTTTTCTTTAAAGTGGTTAGTAACTATTAATCTAGTGTT		2300
CAATCCATTTCCTCAATGTGATTTGTTCACTTACATCTGTTTACGTTATG		2350
CTCTTCTCAGGGGAAATTTTATGAGCTGTATGAGCTAGATGCGGAATTAG		2400
GTCACAAGGAGCTTGACTGGAAGATGACCATGAGTGGTGTGGGAAAATGC		2450
AGACAGGTAAATTAGTTGAAACAACTGGCCTGCTTGAATTATTGTGTCTA		2500
TAAATTTTGACACCACCTTTTGTTTCAGGTTGGTATCTCTGAAAGTGGGA		2550
TAGATGAGGCAGTGCAAAAGCTATTAGCTCGTGGGTAAGGGAACCATCAT		2600
ACTTTATGGAATTCGTTTACTGCTACTTCGGCTAGGATTTAAGAAATGGA		2650
AATCACTTCAAGCATCATTAGTTAGGATCCTGAGAACTCAGGATGTTTTC		2700
TTATTCGTTATAATAAGTCTTTTCATCAAGGAGTAACAAACA		2750
GCACAATATTTGTGTGCTCACTGGCAAGGCATATATACCCAGCTAACCTT		2800
TGCTAGTTCACTGTAGTAACAGTTACGGATAATATATGTTTACTTGTATG		2850
TGGTACCCTCATTTTGTCTCTCATGGAGGCTTTCAAGCCTTGTGTTGAAA		2900
CTGGATAGTTACATATGCTTCCAACAGAAACTAGCATGCAGATTCATATG		2950
CTTTCCTATTCTACTAATTATGTATTGACACACTCGTTGTTTCTTTTGAA	,	3000
AGATATAAAGTTGGACGAATCGAGCAGCTAGAAACATCTGACCAAGCAAA	· .5	3050
AGCCAGAGGTGCTAATACTGTAAGTTTTCTTGGATAGGTCAAGGAGAGTG		3100
TTGCAGACTGTTTTTGATCATTTCTTTTTTTTTTTACATTACTTTCATGCTG	•	3150
TAATTAACTCAATGGCTATTCTGGTCTGATTATCAGATAATTCCAAGGAA		3200
GCTAGTTCAGGTATTAACTCCATCAACAGCAAGCGAGGGAAACATCGGGC		3250
CTGATGCCGTCCATCTTCTTGCTATAAAAGAGGTTTGTTATTTACTTATT		3300
TATCTTATCATGTTCAGTTCATCCAAGTCCTGAAAAATTACACTCTTCTT		3350
TACCAATCTTCCATCAAGCTGTGTAAAGGATTTGGAATTAGAAAATCATT		3400
ATTTGATGCTTTGTTTTATATGCAAGAGGTTCCCTTGAAAAGATCTGTTT		3450
AAGATTCTTTGCACTTGAAAAATTCAATCTTTTTAAGTGAATCCCCTACT		3500
TTCTTACAATGATCATAGTCTGCAATTGCATGTCAAGTAATATCATTCCT		3550
TGTTACTGCATCCCCCTCTTTCTTAATGACCATTGTCTATGTTGTTTTG		3600
TCTCGTGTGCTGGAGAAAATGATAGCTGATCCAAGCTGTACATTATCATG		3650
ATTAAGTAGCTGCTCAGGAATTGCCTTTGGTTACATTGCCTAATGGTTTG		3700
ATGTCAATTTTTCTTCTGAATCTTTATTTTAGATCAAAATGGAGCTACAA		3750
AAGTGTTCAACTGTGTATGGATTTGCTTTTGTTGACTGTGCTGCCTTGAG		3800
GTTTTGGGTTGGGTCCATCAGCGATGATGCATCATGTGCTGCTCTTGGAG		3850
CGTTATTGATGCAGGTAAGCAAGTGTATTCTGTATCTTATGTGTACCATG		3900
TGACTTCCTGTGCATATATTTGGGTTGCAGGAACTAATTCTGAATCACCA		3950
TTTGGTATGTTTTTCCAGGTTTCTCCAAAGGAAGTGTTATATGACAGTA		4000
AAGGTAAACTGCTTGTATCGCCAGTTGTTTTGTTAAACAGAATTTAAGGT		4050
AAATGACACTGGTTAATTTAAAGTGCATACATGTTGAAATATTGCAGGGC		4100
TATCAAGAGAAGCACAAAAGGCTCTAAGGAAATATACGTTGACAGGTACC		4150
ATTTCAGTAGGCAAGCTAACTGACAATTTAACCGCTCACCGAATGATAGG		4200
TCTCTTAAACATTGCTAATGTAGATGATGTTTATGTTTCAATCTAATAGG		4250
GTCTACGGCGGTACAGTTGGCTCCAGTACCACAAGTAATGGGGGATACAG		4300
ATGCTGCTGGAGTTAGAAATATAATAGAATCTAACGGATACTTTAAAGGT		4350
TCTTCTGAATCATGGAACTGTGCTGTTGATGGTCTAAATGAATG		4400

Figure 11 (Continued)

TGCCCTTAGTGCTCTTGGAGAGCTAATTAATCATCTGTCTAGGCTAAAGG	4450
TGTGTTGGCTTGTTTAGTTTTTGCTTTTCACAAATTAAGCAAAGGAACTT	4500
TTCATAACTTACAGTTTCTATCTACTTGCAGCTAGAAGATGTACTTAAGC	4550
ATGGGGATATTTTTCCATACCAAGTTTACAGGGGTTGTCTCAGAATTGAT	4600
GGCCAGACGATGGTAAATCTTGAGATATTTAACAATAGCTGTGATGGTGG	4650
TCCTTCAGGCAAGTGCATATTTCTTTTTTGATAACTTCAACTAGAGGGCA	4700
GACATAGAAGGAAAAATTCTAATACTTCGTACGGATCTCCAGTAAGTA	4750
AGCCGATTTTTGTTTACCTATGTAGGGACCTTGTACAAATATCTTGATAA	4800
CTGTGTTAGTCCAACTGGTAAGCGACTCTTAAGGAATTGGATCTGCCATC	4850
CACTCAAAGATGTAGAAAGCATCAATAAACGGCTTGATGTAGTTGAAGAA	4900
TTCACGGCAAACTCAGAAAGTATGCAAATCACTGGCCAGTATCTCCACAA	4950
ACTTCCAGACTTAGAAAGACTGCTCGGACGCATCAAGTCTAGCGTTCGAT	5000
CATCAGCCTCTGTGTTGCCTGCTCTTCTGGGGAAAAAAGTGCTGAAACAA	5050
CGAGTAAGTATCAATCACAAGTTTTCTGAGTAATGCCTTCCATGAGTAGT	5100
ATAGGACTAAAACATTACGGGTCTAGCTAAAGACTGTTCTCCTTCTTTTG	. 5150
CAATGTCTGGTTATTCATTACATTTCTCTTAACTTATTGCATTGCAGGTT	5200
AAAGCATTTGGGCAAATTGTGAAAGGGTTCAGAAGTGGAATTGATCTGTT	5250
GTTGGCTCTACAGAAGGAATCAAATATGATGAGTTTGCTTTATAAACTCT	5300
GTAAACTTCCTATATTAGTAGGAAAAAGCGGGCTAGAGTTATTTCTTTC	5350
CAATTCGAAGCAGCCATAGATAGCGACTTTCCAAATTATCAGGTGCCCAT	5400
CTATCTTTCATACTTTACAACAAAATGTCTGTCACTACTCAAAGCAATGC	5450
ATATGGCTTAGATCTCAACTCACACCCCGAGGATCCTAAAGGGATTTGCT	5500
TTTTATTCCTAATGTTTTTGGATGGTTTGATTTATTTCTAACTTGAACTT	5550
ATTAATCTTGTACCAGAACCAAGATGTGACAGATGAAAACGCTGAAACTC	5600
TCACAATACTTATCGAACTTTTTATCGAAAGAGCAACTCAATGGTCTGAG	5650
GTCATTCACACCATAAGCTGCCTAGATGTCCTGAGATCTTTTGCAATCGC	5700
AGCAAGTCTCTCTGCTGGAAGCATGGCCAGGCCTGTTATTTTTCCCGAAT	5750
CAGAAGCTACAGATCAGAATCAGAAAACAAAAGGGCCAATACTTAAAATC	5800
CAAGGACTATGGCATCCATTTGCAGTTGCAGCCGATGGTCAATTGCCTGT	5850
TCCGAATGATATACTCCTTGGCGAGGCTAGAAGAAGCAGTGGCAGCATTC	5900
ATCCTCGGTCATTGTTACTGACGGGACCAAACATGGGCGGAAAATCAACT	5950
CTTCTTCGTGCAACATGTCTGGCCGTTATCTTTGCCCAAGTTTGTATACT	6000
CGTTAGATAATTACTCTATTCTTTGCAATCAGTTCTTCAACATGAATAAT	6050
AAATTCTGTTTTCTGTCTGCAGCTTGGCTGCTACGTGCCGTGTGAGTCTT	6100
GCGAAATCTCCCTCGTGGATACTATCTTCACAAGGCTTGGCGCATCTGAT	6150
AGAATCATGACAGGAGAGAGTAAGTTTTGTTCTCAAAATACCAATTCCTC	6200
GAACTATTTACTCAGATTTTGTCTGATTGGACAAGGTGGTTTTGCTTTTT	6250
TTTAGGTACCTTTTTGGTAGAATGCACTGAGACAGCGTCAGTTCTTCAGA	6300
ATGCAACTCAGGATTCACTAGTAATCCTTGACGAACTGGGCAGAGGAACT	6350
AGTACTTTCGATGGATACGCCATTGCATACTCGGTAACCTGCTCTTCTCC	6400
TTCAACTTATACTTGTTGATCAACAAAAACATGCAATTCATTTTGCTGAA	6450
ACTTATTGATTTATATCAGGTTTTTCGTCACCTGGTAGAGAAAGTTCAAT	6500
GTCGGATGCTCTTTGCAACACATTACCACCCTCTCACCAAGGAATTCGCG	6550
TCTCACCCACGTGTCACCTCGA TACACATGGCTTGCGCATTCAAATCAAG	6600
Figure 11 (Continued)	

Figure 11 (Continued)

ATCTGATTATCAACCACGTGGTTGTGATCAAGACCTAGTGTTCTTGTACC	6650
GTTTAACCGAGGGAGCTTGTCCTGAGAGCTACGGACTTCAAGTGGCACTC	6700
ATGGCTGGAATACCAAACCAAGTGGTTGAAACAGCATCAGGTGCTGCTCA	6750
AGCCATGAAGAGATCAATTGGGGAAAACTTCAAGTCAAG	6800
CTGAGTTCTCAAGTCTGCATGAAGACTGGCTCAAGTCATTGGTGGGTATT	6850
TCTCGAGTCGCCCACAACAATGCCCCCATTGGCGAAGATGACTACGACAC	6900
TTTGTTTTGCTTATGGCATGAGATCAAATCCTCTTACTGTGTTCCCAAAT	6950
AAATGGCTATGACATAACACTATCTGAAGCTCGTTAAGTCTTTTGCTTCT	7000
CTGATGTTTATTCCTCTTAAAAAATGCTTATATATCAAAAAATTGTTTCC	7050
TCGATTATAACAAGATTATATATGTATCTGTCGGTTTAGCTATGGTATAT	7100
AATATATGTATGTTCATGAGATTGGTCAAGAGAAAATACTCACAAACAGTA	7150
TATTAAGAAGGAAATATGTTTATGCATTAATTTAAGTTTCAAGATAAACT	. 7200
GCAAATAACCTCGACTAAAGTTGCAAAGACCAAACACAAATTACAAAACT	7250
TATAAGACTTAAGTTCTGAATTCCCTAAAACCAAAAAAAA	. 7300
TATTTTGTTGCATCTACAAACAACACAAACCTACATAGTTTATAACTTAC	7350
TCATCACTGAGATTAACATCAGAATCATTCTCCATTTCTTCATCTTCACT	7400
CTCATCATCACCACCACCATGATGATTCTCCTCCTCTTCACGTAACC	7450
TAGCAATCTCACTCTGAGCTCTATCAACAATCTGCTTCTTCTGCAACTCC	7500
AAATCTCTCTGAAAATCAGCTCTCATCTTCTCCAACTCCTTCATTTGCTC	7550
TTTCTTACTCTTCTCCATCTTCTCATAAACCTTCCCAAACCTCTCAACAG	7600
AATCCGCCAACATCTTATACGAAGCAGCGTCATTAACCTTCTTCCTCTCG	7650
TACTCAACCTCATCCTCATCCTCCTCCTCTCTCAGAATCACCAGGACT	7700
ATCCATCATCTCATCAAACCCATTAGACTTATCTAAATAAA	7750
TCATAAACACAAACTCACCTGAATCAACACCACAAGCTAAACCTAAATCC	7800
GACTTGGGCGAAACACAAAGCAACATATCCAACTTATTGAAAAACGACCA	7850
TTTACTTGAACCTAAACCTGATTTCTCAACCTTAATCTTCTCTTTTCTAT	7900
ACTTCCTCTTCAAGTCATCAATCATTCTCCTACATTGCGTCTCAGATTTC	7950
TCCATCCTTAGCTCCTCACTCACTTTCTCAGCTACTTCATTCCAATCCTC	8000
GTTCCTCAAACTCCTTCTACCCAATTGCAAAAACCTATCTCCCCAAACTT	8050
CAAGCAACACAA	8062

Figure 11 (Continued)




Figure 12

Comments/References: AtMSH3 3' side antisense : AtMSH3 3' (13 = 2104bp) from

pUC18/13 Sal1/Sst1/T4 into pCW164 BamH1/T4 in Agrobacterium LBA44O4




Figure 13

Comments/References: AtMSH6 (S8) 3' side antisens : 62 Sal1/Sst1/T4 (1379bp) into pCW164 BamH1/T4




Figure 14

Comments/References: AtMSH6 3'/AtMSH3 3' antisense: AtMSH6 (S8) 3' side (62=1379bp) Sal1/Sst1/T4 into pPF13 (pCW164 AtMSH3 (S5) 3' side (13=2104) antisens)/Sma1. in LBA4404




Figure 15

Comments/References: AtMSH3 3'/AtMSH6 3' antisens (D): AtMSH3 (S5) 3' side (13=2104bp) Sal1/Sst1/T4 into pPF14 (AtMSH6 (S8) 3'side (62=1379bp) antisense into pCW164)/Sma1. in LBA4404




Figure 16

Comments/References: AtMSH3 (SB) complete, sense orientation : pPF26 (3342bp)

Sma1 into pCW164 Sma1




Figure 17

Comments/References: pPZP111 with codA EcoR1 cassette in EcoR1 site and Hind3 GUS cassette in Hind3 site. KanR. All genes under Promoter/terminator 35S

Figure 18





Figure 19

p3243



PCT/EP98/06977 WO 99/19492

1

SEQUENCE LISTING

<211>

<212>

DNA

```
Rhone-Poulenc Agro; Betzner, Andreas Stefan; Doutriaux,
<110>
             Marie-Pascale; Freyssinet, Georges; Perez, Pascual.
             Methods for obtaining plant varieties
<120>
             395498C
<130>
             PO9745
<150>
             1997-10-10
<151>
            98
<160>
             1
<210>
<211>
             23
             DNA
<212>
             Artificial sequence
<213>
<220>
             modified_base
<221>
<222>
             11
             Į
<223>
<220>
             modified_base
<221>
             14
<222>
<223>
             Ι
<220>
             modified_base
<221>
             17
<222>
<223>
             Ι
<220>
             Degenerate oligonucleotides UPMU used to isolate AtMSH3 and
<223>
             AtMSH6.
<300>
<301>
             Reenan and Kolodner
             Genetics
<302>
             132
<303>
            963-973
<306>
              1992
 <307>
              1
 <400>
                                                                       23
 ctggatccac nggnccnaay atg
              2
 <210>
              23
```

```
Artificial sequence
<213>
<220>
             modified base
<221>
<222>
             15
<223>
             Ι
<220>
             modified base
<221>
<222>
             18
<223>
             I
<220>
             Degenerate oligonucleotides DOMU used to isolate AtMSH3 and
<223>
             AtMSH6.
<300>
             Reenan and Kolodner
<301>
<302>
             Genetics
<303>
             132
<306>
             963-973
            1992
<307>
             2
<400>
                                                                       23
ctggatccrt artgngtnrc raa
<210>
             3
             24
<211>
<212>
             DNA
             Artificial sequence
<213>
<220>
             MSH3 specific primer 636 for PCR using cDNA of Arabidopsis
<223>
             thaliana ecotype Columbia
             3
<400>
                                                                        24
tgctagtgcc tcttgcaagc tcat
 <210>
              27
<211>
              DNA
<212>
              Artificial sequence
 <213>
 <220>
              Primer AP1 for PCR using cDNA of Arabidopsis thaliana ecotype
 <223>
              Columbia containing adapter sequences ligated to both its
              ends
 <400>
```

ccatcctaat	acgactcact atagggc	27
<210>	5	
<211>	23	
<212>	DNA	
<213>	Artificial sequence	
<213>	nterrora bequess	
<220>		
<223>	Primer AP2 for PCR using cDNA of Arabidopsis thaliand Columbia containing adapter sequences ligated to both ends	a ecotyp h its
<400>	5	
actcactata	a gggctcgagc ggc	23
		•
<210>	6	
<211>	30	
<211>	DNA	
-	Artificial sequence	
<213>	Altificial Sequence	
<220>		
<223>	MSH3 specific primer S525 for PCR using cDNA of Arab thaliana ecotype Columbia	oldobsis
<400>	6	
aggttctgat	t tatgtgtgac gctttactta	30
<210>	7	
<211>	29	
<212>	DNA	
<213>	Artificial sequence	
<220>	The second of Aught	idoneie
<223>	MSH3 specific primer S51 for PCR using cDNA of Arab thaliana ecotype Columbia	Idobala
<400>	7	
ggatcgggt	ta ctgggttttg agtgtgagg	29
<210>	8	
<211>	24	
<212>	DNA	
<213>	Artificial sequence	
	<u>.</u>	
<220>	, niva - E lumb	i donei c
<223>	MSH3 specific primer 635 for PCR using cDNA of Arab	Tachera

4

<400>	8		
gcacgtgctt	gatggtgttt tcac		24
	•		
<210>	9		
<211>	28		
	DNA		
<213>	Artificial sequence		
<220>			
<223>	MSH3 specific primer S523 for PCR thaliana ecotype Columbia	using cDNA of	Arabidopsi
<400>	9		
tcagacagta	tccagcatgg cagaagta		28
		*	•
		•	1
<210>	10	• • • • • • • • • • • • • • • • • • • •	·c,
<211>	33		
<212>	DNA		
<213>	Artificial sequence		
<220>			
<223>	MSH3 specific primer 1S5 for PCR thaliana ecotype Columbia	using cDNA of	Arabidopsis
<400>	10		
atcccgggat	gggcaagcaa aagcagcaga cga		33
210-	11		
<210> <211>	11 27		
<211>	DNA		
	Artificial sequence		
<220>			
<223>	MSH3 specific primer S53 for PCR	using cDNA of	Arabidopsis
(223)	thaliana ecotype Columbia	asing come or	
<400>	11		
gacaaagago	gaaatgagge eeettgg		27
<210>	12		
<211>	1250		
<212>	DNA		
<213>	Arabidopsis thaliana ecotype Col	umbia	
<223>	Clone 52	<del></del>	

<400> 12

cccgggatgg	gcaagcaaaa	gcagcagacg	atttctcgtt	tcttcgctcc	caaacccaaa	60
tccccgactc	acgaaccgaa	tccggtagcc	gaatcatcaa	caccgccacc	gaagatatcc	120
gccactgtat	ccttctctcc	ttccaagcgt	aagcttctct	ccgaccacct	cgccgccgcg	180
tcacccaaaa	agcctaaact	ttctcctcac	actcaaaacc	cagtacccga	tcccaattta	240
caccaaagat	ttctccagag	atttctggaa	ccctcgccgg	aggaatatgt	tcccgaaacg	300
tcatcatcga	ggaaatacac	accattggaa	cagcaagtgg	tggagctaaa	gagcaagtac	360
ccagatgtgg	ttttgatggt	ggaagttggt	tacaggtaca	gattcttcgg	agaagacgcg	420
gagatcgcag	cacgcgtgtt	gggtatttac	gctcatatgg	atcacaattt	catgacggcg	480
agtgtgccaa	catttcgatt	gaatttccat	gt <b>gagaa</b> gac	tggtgaatgc	aggatacaag	540
attggtgtag	tgaagcagac	tgaaactgca	gccattaagt	cccatggtgc	aaaccggacc	600
ggcccttttt	tccggggact	greggegrig	tataccaaag	ccacgcttga	agcggctgag	660
gatataagtg	gtggttgtgg	tggtgaagaa	ggttttggtt	cacagagtaa	tttcttggtt	720
tgtgttgtgg	atgagagagt	taagtcggag	acattaggct	gtggtattga	aatgagtttt	780
gatgttagag	teggtgttgt	tggcgttgaa	atttcgacag	gtgaagttgt	ttatgaagag	840
ttcaatgata	atttcatgag	aagtggatta	gaggctgtga	ttttgagctt	gtcaccagct	900
gagctgttgc	ttggccagcc	tctttcacaa	caaactgaga	agtttttggt	ggcacatgct	960
ggacctacct	caaacgttcg	agtggaacgt	gcctcactgg	attgtttcag	caatggtaat	1020
gcagtagatg	aggttatttc	attatgtgaa	aaaatcagcg	caggtaactt	agaagatgat	1080
aaagaaatga	agctggaggc	tgctgaaaaa	ggaatgtctt	gcttgacagt	tcatacaatt	1140
atgaacatgc	cacatctgac	tgttcaagcc	ctcgccctaa	cgttttgcca	tctcaaacag	1200
tttggatttg	aaaggatcct	ttaccaaggg	gcctcatttc	gctctttgtc		1250

<sup>&</sup>lt;210> 13

<sup>&</sup>lt;211> 34

<sup>&</sup>lt;212> DNA

<sup>&</sup>lt;213> Artificial sequence

<sup>&</sup>lt;220>

<sup>&</sup>lt;223> MSH3 specific primer 2S5 for PCR using cDNA of Arabidopsis chaliana ecotype Columbia

<400>	13					
atcccgggtc	aaaatgaaca	agttggtttt	agtc			34
<210>	14					
<211>	27					
<212>	DNA					
<213>	Artificia	l seguence				
(213)						
<220>						
<223>		ific primer ecotype Col		R using cDN	A of Arabido	psis
<400>	14					
gccacatctg	actgttcaag	ccctcgc			•	27
5	<u> </u>	_		•		
					•	
<210>	15			•	į	
<211>	2110				•	
<212>	DNA		_			
	Arabidops	is thailana	ecotype Co	lumbla		
<223>	Clone 13					
<400>	15					
gccacatctg	actgttcaag	ccctcgccct	aacgttttgc	catctcaaac	agtttggatt	60
tgaaaggatc	ctttaccaag	gggcctcatt	tcgctctttg	tcaagtaaca	cagagatgac	120
teteteagee	aatactctgc	aacagttgga	ggttgtgaaa	aataattcag	atggatcgga	180
atctggctcc	ttattccata	atatgaatca	cacacttaca	gtatatggtt	ccaggcttct	240
tagacactgg	gtgactcatc	ctctatgcga	tagaaatttg	atatctgctc	ggcttgatgc	300
tgtttctgag	atttctgctt	gcatgggatc	tcatagttct	tcccagctca	gcagtgagtt	360
ggttgaagaa	ggttctgaga	gagcaattgt	atcacctgag	ttttatctcg	tgctctcctc	420
agtcttgaca	gctatgtcta	gatcatctga	tattcaacgt	ggaataacaa	gaatctttca	480
teggaetget	aaagccacag	agttcattgc	agttatggaa	gctattttac	ttgcggggaa	540
gcaaattcag	cggcttggca	taaagcaaga	ctctgaaatg	aggagtatgc	aatctgcaac	600
tgtgcgatct	actcttttga	gaaaattgat	ttctgttatt	tcatcccctg	ttgtggttga	660
caatgccgga	aaacttctct	ctgccctaaa	taaggaagcg	gctgttcgag	gtgacttgct	720
cgacatacta	atcacttcca	gcgaccaatt	teetgagett	gctgaagctc	gccaagcagt	780
rttagtcatc	agggaaaagc	tggattcctc	gatagcttca	tttcgcaaga	agctcgctat	P40

togaaatttg	gaatttcttc	aagtgtcggg	gatcacacat	ttgatagagc	tgcccgttga	900
ttccaaggtc	cctatgaatt	gggtgaaagt	aaatagcacc	aagaagacta	ttcgatatca	960
tcccccagaa	atagtagctg	gcttggatga	gctagctcta	gcaactgaac	atcttgccat	1020
tgtgaaccga	gcttcgtggg	atagtttcct	caagagtttc	agtagatact	acacagattt	1080
taaggctgcc	gttcaagctc	ttgctgcact	ggactgtttg	cactcccttt	caactctatc	1140
tagaaacaag	aactatgtcc	gtcccgagtt	tgtggatgac	tgtgaaccag	ttgagataaa	1200
catacagtct	ggtcgtcatc	ctgtactgga	gactatatta	caagataact	tcgtcccaaa	1260
tgacacaatt	ttgcatgcag	aaggggaata	ttgccaaatt	atcaccggac	ctaacatggg	1320
aggaaagagc	tgctatatcc	gtcaagttgc	tttaatttcc	ataatggctc	aggttggttc	1380
ctttgtacca	gcgtcattcg	ccaagctgca	cgtgcttgat	ggtgttttca	cccggacggg	1440
tgcttcagac	agtatccagc	atggcagaag	tacctttcta	gaagaattaa	gtgaagcgtc	1500
acacataatc	agaacctgtt	cttctcgttc	gcttgttata	ttagatgagc	ttggaagagg	1560
cactagcaca	cacgacggtg	tagccattgc	ctatgcaaca	ttacagcatc	tcctagcaga	1620
aaagagatgt	ttggttcttt	ttgtcacgca	ttaccctgaa	atagctgaga	tcagtaacgg	1680
attcccaggt	tctgttggga	cataccatgt	ctcgtatctg	acattgcaga	aggataaagg	1740
cagttatgat	catgatgatg	tgacctacct	atataagctt	gtgcgtggtc	tttgcagcag	1800
gagctttggt	tttaaggttg	ctcagcttgc	ccagatacct	ccatcatgta	tacgtcgagc	1860
catttcaatg	gctgcaaaat	tggaagctga	ggtacgtgca	agagagagaa	atacacgcat	1920
gggagaacca	gaaggacatg	aagaaccgag	aggcgcagaa	gaatctattt	cggctctagg	1980
tgacttgttt	gcagacctga	aatttgctct	ctctgaagag	gacccttgga	aagcattcga	2040
gtttttaaag	catgcttgga	agattgctgg	caaaatcaga	ctaaaaccaa	cttgttcatt	2100
ttgacccggg						2110

<210>	.16
<211>	29
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	MSH3 specific primer S51 for PCR u

MSH3 specific primer S51 for PCR using cDNA of Arabidopsis thaliana ecotype Columbia

<400> 16

ggat	eggg	ca c	tggg	TTTT	g ag	tgtg	agg										29
<210:	>		17														
<211:			30														
<212			DNA														
<213:			Artificial sequence														
(413)	•		AIC.	1110	141 .	ocqu.											
<220:					<b>.</b> .	:		- CF	ar 6.	D	CD	-:	~DM	λ of	A ~ = b	idone	ic
<223:	>			_	a ec					OI P	CR U	sing	CDN.	A OL	ALGL	idops	
<400	>		17														
aggttctgat tatgtgtgac gctttactta 30														30			
<210:	>		18														
<211	>		352	2													
<212	>		DNA														
<213	>		Ara	bido	psis	tha	lian	a ec	otyp	e Co	lumb	ia	,			· e.	
<220	>																
<221	>		CDS														
<222			(10	٥)	(3	342)											
<223							ath	cDNA	and	ded	uced	sea	uenc	e of	the	encod	ed
~223					tide		J					•					
			por	, pop													
<400	>		18														
					_ ~						F 2 6 6		, , , ,		CCEE	_	60
ccta	agaa	ag c	gege	gaaa	ia cc	.ggca	accc	aag	reeg	icca	cage	cacy	ac c	acya	cctt	-	00
catt	tctc	tt a	aaacg	gagg	ra ga	ttac	gaat	aaa	ıgcaa	ict							99
atq	aac	aaq	caa	aaq	cag	cag	acg	att	tct	cgt	ttc	ttc	gct	CCC	aaa	1	47
Met	Glv	Lvs	Gln	Lvs	Gln	Gln	Thr	Ile	Ser	Arg	Phe	Phe	Ala	Pro	Lys		
1		- 4		15					10	_				15			
•				_													
CCC	222	tcc	ccg	act	cac	gaa	cca	aat	cca	αta	acc	gaa	tca	tca	aca	1	.95
			Pro														
PLO	ב עם	Jer	20	1111	1110	010		25		• • • •			30				
			20					23					50				
			aag			~~~	205	~= >	F 00	++0	+ ~+	cct	tcc	aan	cat	9	243
																_	
Pro	Pro		Lys	Tie	ser	Ala		vai	ser	Pne	ser		ser	ьys	ALG		
		35					40					45					
-	•			•												_	
			tcc													2	291
Lys	Leu	Leu	Ser	Asp	His	Leu	Ala	Ala	Ala	Ser		Lys	Lys	Pro	ьуs		
	50					55					60						
			cac													3	339
Leu	Ser	Pro	His	Thr	Gln	Asn	Pro	Val	Pro	Asp	Pro	Asn	Leu	His	Gln		
65					70					75					80		

								ccc Pro								387
								aca Thr 105								435
_							-	gtg Val	_	_	_		_	_		483
								gac Asp								531
								cac His							gtg Val 160	 579
								gtg Val							gga Gly	627
								act Thr 185			_	_		_		675
								ttt Phe								723
								gct Ala								771
ggt Gly 225	ggt Gly	gaa Glu	gaa Glu	ggt Gly	ttt Phe 230	ggt Gly	tca Ser	cag Gln	agt Ser	aat Asn 235	ttc Phe	ttg Leu	gtt Val	cya	gtt Val 240	819
gtg Val	gat Asp	gag Glu	aga Arg	gtt Val 245	aag Lys	tcg Ser	gag Glu	aca Thr	tta Leu 250	ggc	tgt Cys	ggt Gly	att Ile	gaa Glu 255	atg Met	867
								gtt Val 265								915
								gat <b>As</b> p								963

			att Ile													10:	11
			caa Gln													109	59
			gtt Val													110	)7
			gta Val 340													115	55
			gaa Glu												aaa Lys	120	)3
			tgc Cys												ctg 🧍 Leu	12!	51
			gcc Ala													12:	99
ttt Phe	gaa Glu	agg Arg	atc Ile	ctt Leu 405	tac Tyr	caa Gln	Gly	gcc Ala	tca Ser 410	Phe	cgc Arg	tct Ser	ttg Leu	tca Ser 415	agt Ser	13	47
			atg Met 420													13	95
			aat Asn					Glu								14	43
		His	aca Thr				Tyr					Leu				14	91
	Thr		cct Pro			Asp					Ser					15	39
			gag Glu		Ser					/ Ser					Gln	1.5	87

	_	-		_			-				_	_	att Ile 510		1635
									_	_			atg Met		1683
		_			_				_				cgg Arg		1731
													ctt Leu		1779
•			-					_		_		_	atg Met	 	1827
_													ttg Leu 590	tct. Ser	1875
_													ctt Leu		1923
_			_	_		_	_	_					gac Asp		1971
													cgc Arg		2019
													tca Ser		2067
													tcg Ser 670		2115
			Ile		_		-	-		_			atg His		2163
		Val										His	ccc Pro		2211

								ctt Leu		2259
								agt Ser 735		2307
								ctg Leu		2355
								gtc Val		2403
								cag Gln		2451
								gtc Val	.5	2499
								atc Ile 815		2547
								gct Ala		2595
								ttc Phe		2643
								tca Ser		2691
_	_		_	_				gaa Glu		2739
			Thr					tta Leu 895		2787
		Gly						gcc Ala		2835

gca aca tta cag cat ctc cta gca gaa aag aga tgt ttg gtt ctt ttt Ala Thr Leu Gln His Leu Leu Ala Glu Lys Arg Cys Leu Val Leu Phe 915 920 925	2883
gtc acg cat tac cct gaa ata gct gag atc agt aac gga ttc cca ggt Val Thr His Tyr Pro Glu Ile Ala Glu Ile Ser Asn Gly Phe Pro Gly 930 935 940	2931
tct gtt ggg aca tac cat gtc tcg tat ctg aca ttg cag aag gat aaa Ser Val Gly Thr Tyr His Val Ser Tyr Leu Thr Leu Gln Lys Asp Lys 945 950 955 960	2979
ggc agt tat gat cat gat gtg acc tac cta tat aag ctt gtg cgt Gly Ser Tyr Asp His Asp Asp Val Thr Tyr Leu Tyr Lys Leu Val Arg 965 970 975	3027
ggt ctt tgc agc agg agc ttt ggt ttt aag gtt gct cag ctt gcc cag Gly Leu Cys Ser Arg Ser Phe Gly Phe Lys Val Ala Gln Leu Ala Gln 980 985 990	3075
ata cct cca tca tgt ata cgt cga gcc att tca atg gct gca aaa ttg .  Ile Pro Pro Ser Cys Ile Arg Arg Ala Ile Ser Met Ala Ala Lys Leu 995 1000 1005	3123
gaa get gag gta egt gea aga gag aga aat aca ege atg gga gaa eea Glu Ala Glu Val Arg Ala Arg Glu Arg Asn Thr Arg Met Gly Glu Pro 1010 1015 1020	3171
gaa gga cat gaa gaa ccg aga ggc gca gaa gaa tct att tcg gct cta Glu Gly His Glu Glu Pro Arg Gly Ala Glu Glu Ser Ile Ser Ala Leu 1025 1030 1035 1040	3219
ggt gac ttg ttt gca gac ctg aaa ttt gct ctc tct gaa gag gac cct Gly Asp Leu Phe Ala Asp Leu Lys Phe Ala Leu Ser Glu Glu Asp Pro 1045 1050 1055	3267
tgg aaa gca ttc gag ttt tta aag cat gct tgg aag att gct ggc aaa Trp Lys Ala Phe Glu Phe Leu Lys His Ala Trp Lys Ile Ala Gly Lys 1060 1065 1070	3315
atc aga cta aaa cca act tgt tca ttt tgatttaatc ttaacattat Ile Arg Leu Lys Pro Thr Cys Ser Phe 1075 1080	3362
agcaactgca aggtettgat catetgttag ttgegtaeta aettatgtgt attagtataa	3422
caagaaaaga gaattagaga gatggattet aateeggtgt tgeagtaeat etttteteea	3482
cccgcataaa aaaaaaaaa aaaaaaaaaaaaaaaa	3522
210> 19	

<210> 19 <211> .081 <212> PRT

<21:					opsi: ptide			na eo	coty	pe Co	olumi	oia			
< 400	)>		19												
Met 1	Gly	Lys	Gln	Lys 5	Gln	Gln	Thr	Ile	Ser 10	Arg	Phe	Phe	Ala	Pro 15	Lys
Pro	Lys	Ser	Pro 20	Thr	His	Glu	Pro	Asn 25	Pro	Val	Ala	Glu	Ser 30	Ser	Thr
Pro	Pro	Pro 35	Lys	Ile	Ser	Ala	Thr 40	Val	Ser	Phe	Ser	Pro 45	Ser	Lys	Arg
Lys	Leu 50	Leu	Ser	Asp	His	Leu 55	Ala	Ala	Ala	Ser	Pro 60	Lys	Lys	Pro	Lys
Leu 65	Ser	Pro	His	Thr	Gln 70	Asn	Pro	Val	Pro	Asp 75	Pro	Asn	Leu	His	Gln 80
Arg	Phe	Leu	Gln	Arg 85	Phe	Leu	Glu	Pro	Ser 90	Pro	Glu	Glu	Tyr	Val 95	Pro
Glu	Thr	Ser	Ser 100	Ser	Arg	Lys	Tyr	Thr 105	Pro	Leu	Glu	Gln	Gln 110	Val	Val
Glu	Leu	Lys 115	Ser	Lys	Tyr	Pro	Asp 120	Val	Val	Leu	Met	Val 125	Glu	Val	Gly
Tyr	Arg 130	Tyr	Arg	Phe	Phe	Gly 135	Glu	Asp	Ala	Glu	Ile 140	Ala	Ala	Arg	Val
Leu 145	Gly	Ile	Tyr	Ala	His 150	Met	Asp	His	Asn	Phe 155	Met	Thr	Ala	Ser	Val 160
Pro	Thr	Phe	Arg	Leu 165	Asn	Phe	His	Val	Arg 170	Arg	Leu	Val	Asn	Ala 175	Gly
туr	Lys	Ile	Gly 180	Val	Val	Lys	Gln	Thr 185	Glu	Thr	Ala	Ala	Ile 190	Lys	Ser
His	Gly	Ala 195	Asn	Arg	Thr	Gly	Pro 200	Phe	Phe	Arg	Gly	Leu 205	Ser	Ala	Leu
Tyr	Thr 210	Lys	Ala	Thr	Leu	Glu 215	Ala	Ala	Glu	Asp	Ile 220	Ser	Gly	Gly	Cys
Gly 225	Gly	Glu	Glu	Gly	Phe 230	Gly	Ser	Gln	Ser	Asn 235	Phe	Leu	Val	Cys	Val 240
Val	Asp	Glu	Arg	Val 245	_	Ser	Glu	Thr	Leu 250	Gly	Cys	Gly	Ile	Glu 255	Met

Ser	Phe	Asp	Val 260	Arg	Val	Gly	Val	Val 265	Gly	Val	Glu	Ile	Ser 270	Thr	Gly
Glu	Val	Val 275	Tyr	Glu	Glu	Phe	Asn 280	Asp	Asn	Phe	Met	Arg 285	Ser	Gly	Leu
Glu	Ala 290	Val	Ile	Leu	Ser	Leu 295	Ser	Pro	Ala	Glu	Leu 300	Leu	Leu	Gly	Gln
Pro 305	Leu	Ser	Gln	Gln	Thr 310	Glu	Lys	Phe	Leu	Val 315	Ala	Met	Ala	Gly	Pro 320
Thr	Ser	Asn	Val	Arg 325	Val	Glu	Arg	Ala	Ser 330	Leu	Asp	Суз	Phe	Ser 335	Asn
-			340	_				345			Glu		350		
•		355		_			360				Glu	365			
•	370		·			375					Asn 380		٠		
385					390					395	Leu				400
				405			-		410		Arg			415	
			420					425			Gln		430		
		435					440				Ser	445			
	450					455					Leu 460				
465					470					475					480
				485					490	1	His			495	
			<b>50</b> 0					505			Arg		510		
		515	;				520	•			Thr	525			
Ser	Ser 520	_	) Ile	Gln	Arg	Gly ברכ		rnr	Arg	, ile	Phe 540		arg	inr	ATS

	Lys 545	Ala	Thr	Glu	Phe	Ile 550	Ala	Val	Met	Glu	Ala 555	Ile	Leu	Leu	Ala	Gly 560
1	ŗķs	Gln	Ile	Gln	Arg 565	Leu	Gly	Ile	Lys	Gln 570	Asp	Ser	Glu	Met	Arg 575	Ser
ı	let	Gln	Ser	Ala 580	Thr	Val	Arg	Ser	Thr 585	Leu	Leu	Arg	Lys	Leu 590	Ile	Ser
1	/al	Ile	Ser 595	Ser	Pro	Val	Val	Val 600	Asp	Asn	Ala	Gly	Lys 605	Leu	Leu	Ser
7	Ala	Leu 610	Asn	Lys	Glu	Ala	Ala 615	Val	Arg	Gly	Asp	Leu 620	Leu	Asp	Ile	Leu
	(le 525	Thr	Ser	Ser	Asp	Gln 630	Phe	Pro	Glu	Leu	Ala 635	Glu	Ala	Arg	Gln	Ala 640
,	/al	Leu	Val	Ile	Arg 645	Glu	Lys	Leu	Asp	Ser 650	Ser	Ile	Ala	Ser	Phe 655	_
I	Lys	Lys	Leu	Ala 660	Ile	Arg	Asn	Leu	Glu 665	Phe	Leu	Gln	Val	Ser 670	Gly	Ile
7	Chr	His	Leu 675	Ile	Glu	Leu	Pro	Val 680	Asp	Ser	Lys	Val	Pro 685	His	Asn	Trp
•	/al	Lys 690	Val	Asn	Ser	Thr	Lys 695	Lys	Thr	Ile	Arg	Tyr 700	His	Pro	Pro	Glu
	11e 705	Val	Ala	Gly	Leu	Asp 710	Glu	Leu	Ala	Leu	Ala 715	Thr	Glu	His	Leu	Ala 720
•	Ile	Val	Asn	Arg	Ala 725	Ser	Trp	qzA	Ser	Phe 730	Leu	Lys	Ser	Phe	Ser 735	Arg
•	Tyr	Tyr	Thr	Asp 740	Phe	Lys	Ala	Ala	Val 745		Ala	Leu	Ala	Ala 750	Leu	Asp
•	Cys	Leu	His 755	Ser	Leu	Ser	Thr	Leu 760	Ser	Arg	Asn	Lys	Asn 765	Tyr	Val	Arg
	Pro	Glu 770	Phe	Val	Asp	Asp	Cys 775	Glu	Pro	Val	Glu	Ile 780	Asn	Ile	Gln	Ser
	Gly 785	Arg	His	Pro	Val	Leu 790	Glu	Thr	Ile	Leu	Gln 795	Asp	Asn	Phe	Val	Pro 800
	Asn	Asp	Thr	Ile	Leu 805	His	Ala	Glu	Gly	Glu 810		Cys	Gln	Ile	Ile 815	Thr
	Gly	Pro	Asn	Met 820		Gly	Lys	Ser	Cys	_	Ile	Arg	Gln	Val 830		Leu

- Ile Ser Ile Met Ala Gln Val Gly Ser Phe Val Pro Ala Ser Phe Ala 835 840 845
- Lys Leu His Val Leu Asp Gly Val Phe Thr Arg Met Gly Ala Ser Asp 850 860
- Ser Ile Gln His Gly Arg Ser Thr Phe Leu Glu Glu Leu Ser Glu Ala 865 870 875 880
- Ser His Ile Ile Arg Thr Cys Ser Ser Arg Ser Leu Val Ile Leu Asp 885 890 895
- Glu Leu Gly Arg Gly Thr Ser Thr His Asp Gly Val Ala Ile Ala Tyr 900 905 910
- Ala Thr Leu Gln His Leu Leu Ala Glu Lys Arg Cys Leu Val Leu Phe 915 920 925
- Val Thr His Tyr Pro Glu Ile Ala Glu Ile Ser Asn Gly Phe Pro Gly 930 935 940
- Ser Val Gly Thr Tyr His Val Ser Tyr Leu Thr Leu Gln Lys Asp Lys 945 950 955 960
- Gly Ser Tyr Asp His Asp Asp Val Thr Tyr Leu Tyr Lys Leu Val Arg 965 970 975
- Gly Leu Cys Ser Arg Ser Phe Gly Phe Lys Val Ala Gln Leu Ala Gln
  980 985 990
- Ile Pro Pro Ser Cys Ile Arg Arg Ala Ile Ser Met Ala Ala Lys Leu 995 1000 1005
- Glu Ala Glu Val Arg Ala Arg Glu Arg Asn Thr Arg Met Gly Glu Pro 1010 1015 1020
- Glu Gly His Glu Glu Pro Arg Gly Ala Glu Glu Ser Ile Ser Ala Leu 1025 1030 1035 1040
- Gly Asp Leu Phe Ala Asp Leu Lys Phe Ala Leu Ser Glu Glu Asp Pro 1045 1050 1055
- Trp Lys Ala Phe Glu Phe Leu Lys His Ala Trp Lys Ile Ala Gly Lys 1060 1065 1070
- Ile Arg Leu Lys Pro Thr Cys Ser Phe 1075 1080
  - <210> 20
  - <211> 24
  - <212> DNA
  - <213> Artificial sequence

WO 99/19492 PCT/EP98/06977

18

<220> <223>	MSH6 specific primer 638 for PCR using cDNA of Arabidopsis thaliana ecotype Columbia	5
<400>	20	
tctctaccag	gtgacgaaaa accg	24
<210>	21	
<211>	28	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Primer S81 for PCR using cDNA of Arabidopsis thaliana ecot Columbia	ype
<400>	21	
cgtcgccttt	agcatcccct tccttcac	28
<210>	22	
<211>	30	
<212>	DNA	
	Artificial sequence	
<220>		
<223>	MSH6 specific primer S823 for PCR using cDNA of Arabidopsi thaliana ecotype Columbia	S
<400>	22	
gcttggcgca	tctaatagaa tcatgacagg	30
<210>	23	
<211>	24	
<212>	DNA	
<213>	Artificial sequence	
<220> ·		
<223>	MSH6 specific primer 637 for PCR using cDNA of Arabidopsis	ı
	thaliana ecotype Columbia	
<400>	23	
gacagegtea	gttcttcaga atgc	24
<210>	24	
<211>	33	

<212> DNA

WO 99/19492

19

<213>	Artificial sequence	
<220>		
	MCUC analisis animan 100 Sau papa at a appropriate	
<223>	MSH6 specific primer 1S8 for PCR using cDNA of Arabidopsi thaliana ecotype Columbia	S
<400>	24	
atcccgggat	gcagcgccag agatcgattt tgt	33
<210>	25	
<211>	27	
<212>	DNA	
<213>	Artificial sequence	
(413)	Attiticial bequence	
<220>		
<223>	MSH6 specific primer S83 for PCR using cDNA of Arabidopsis thaliana ecotype Columbia	5
	·•	
<400>	25	
cgctatctat	ggetgetteg aattgag	27
<210>	26	
<211>	1385	
<212>	DNA	
<213>	Arabidopsis thaliana ecotype Columbia	
<223>	Clone 43	
<400>	26	
cccgggatgc	agegecagag ategattitg tettiettee aaaaacecae ggeggegaet	60
acgaagggtt	tggtttccgg cgatgctgct agcggcgggg gcggcagcgg aggaccacga	120
tttaatgtga	aggaagggga tgctaaaggc gacgcttctg tacgttttgc tgtttcgaaa	180
tctgtcgatg	aggttagagg aacggatact ccaccggaga aggttccgcg tcgtgtcctg	240
ccgtctggat	ttaageegge tgaateegee ggtgatgett egteeetgtt eteeaatatt	300
atgcataagt	ttgtaaaagt cgatgatcga gattgttctg gagagaggag ccgagaagat	360
gttgttccgc	tgaatgattc atctctatgt atgaaggcta atgatgttat tcctcaattt	120
cgttccaata	atggtaaaac tcaagaaaga aaccatgctt ttagtttcag tgggagagct	180
gaacttagat	cagtagaaga tataggagta gatggcgatg ttcctggtcc agaaacacca	540
gggatgcgtc	cacgtgcttc tcgcttgaag cgagttctgg aggatgaaat gacttttaag	500
gaggataagq	ttcctgtatt ggactctaac aaaaqqctga aaatqctcca ggatccqqtt 6	660

tgtggagaga	agaaagaagt	aaacgaagga	accaaatttg	aatggcttga	gtcttctcga	720
atcagggatg	ccaatagaag	acgtcctgat	gatccccttt	acgatagaaa	gaccttacac	780
ataccacctg	atgttttcaa	gaaaatgtct	gcatcacaaa	agcaatattg	gagtgttaag	840
agtgaatata	tggacattgt	gcttttcttt	aaagtgggga	aattttatga	gctgtatgag	900
ctagatgcgg	aattaggtca	caaggagctt	gactggaaga	tgaccatgag	tggtgtggga	960
aaatgcagac	aggttggtat	ctctgaaagt	gggatagatg	aggcagtgca	aaagctatta	1020
gctcgtggat	ataaagttgg	acgaatcgag	cagctagaaa	catctgacca	agcaaaagcc	1080
agaggtgcta	atactataat	tccaaggaag	ctagttcagg	tattaactcc	atcaacagca	1140
agcgagggaa	acatcgggcc	tgatgccgtc	catettettg	ctataaaaga	gatcaaaatg	1200
gagctacaaa	agtgttcaac	tgtgtatgga	tttgcttttg	ttgactgtgc	tgccttgagg	1250
ttttgggttg	ggtccatcag	cgatgatgca	tcatgtgctg	ctcttggagc	gttattgatg	1320
caggittete	caaaggaagt	gttatatgac	agtaaagggc	tatcaagaga	agcacaaaag	1380
gctctaagga	aatatacgtt	gacagggtct	acggcggtac	agttggctcc	agtaccacaa	1440
gtaatggggg	atacagatgc	tgctggagtt	agaaatataa	tagaatctaa	cggatacttt	1500
aaaggttctt	ctgaatcatg	gaactgtgct	gttgatggtc	taaatgaatg	tgatgttgcc	1560
cttagtgctc	ttggagagct	aattaatcat	ctgtctaggc	taaagctaga	agatgtactt	1620
aagcatgggg	atattttcc	ataccaagtt	tacaggggtt	gtctcagaat	tgatggccag	1680
acgatggtaa	atcttgagat	atttaacaat	agctgtgatg	gtggtccttc	agggaccttg	1740
tacaaatato	ttgataactg	tgttagtcca	actggtaagc	gactcttaag	gaattggatc	1800
tgccatccac	: tcaaagatgt	: agaaagcato	aataaacggc	ttgatgtagt	tgaagaattc	1860
acggcaaact	: cagaaagtat	gcaaatcact	ggccagtato	tccacaaact	tccagactta	1920
gaaagactgo	c toggacgcat	caagtctago	gttcgatcat	cagcctctgt	gttgcctgct	1980
cttctgggg	a aaaaagtgc	gaaacaacga	a gttaaagcat	ttgggcaaat	tgtgaaaggg	2040
ttcagaagt	g gaattgatc	t gttgttggci	ctacagaagg	, aatcaaatat	gatgagtttg	2100
ctttataaa	c totgtaaac	t tootatatta	a gtaggaaaaa	a gegggetaga	a gttatttctt	216
tctcaattc	g aagcagcca	t agatagcg				218

<210> <211> <212> <213> <223>	27 1385 DNA Arabidopsis thaliana ecotype Columbia Clone 62	
<400>	27	
catcagcctc	tgtgttgcct gctcttctgg ggaaaaaagt gctgaaacaa cgagttaaag 60	
catttgggca	aattgtgaaa gggttcagaa gtggaattga tctgttgttg gctctacaga 120	
aggaatcaaa	tatgatgagt ttgctttata aactetgtaa aetteetata ttagtaggaa 180	
aaagcgggct	agagttattt ctttctcaat tcgaagcagc catagatagc gactttccaa 240	
attatcagaa	ccaagatgtg acagatgaaa acgctgaaac tctcacaata cttatcgaac 300	
tttttatcga	aagagcaact caatggtotg aggtoattoa caccataago tgootagatg 360	
tcctgagatc	ttttgcaatc gcagcaagtc tetetgetgg aagcatggee aggeetgtta 420	
tttttcccga	atcagaaget acagateaga atcagaaaac aaaagggeea ataettaaaa 480	
tccaaggact	atggcatcca tttgcagttg cagccgatgg tcaattgcct gttccgaatg 540	
atatactcct	tggcgagget agaagaagea gtggcageat teateetegg teattgttae 600	
tgacgggacc	aaacatgggc ggaaaatcaa ctcttcttcg tgcaacatgt ctggccgtta 660	ı
tctttgccca	acttggctgc tacgtgccgt gtgagtcttg cgaaatctcc ctcgtggata 720	I
ctatcttcac	aaggettgge geatetgata gaateatgae aggagagagt acetttttgg 780	i
tagaatgcac	tgagacageg teagttette agaatgeaae teaggattea etagtaatee 840	ı
ttgacgaact	gggcagagga actagtactt tcgatggata cgccattgca tactcggttt 900	)
ttcgtcacct	ggtagagaaa gttcaatgtc ggatgctctt tgcaacacat taccaccctc 960	ı
tcaccaagga	attegegtet cacceaegtg teacetegaa acacatgget tgegeattea 1020	ı
aatcaagatc	tgattatcaa ccacgtggtt gtgatcaaga cctagtgttc ttgtaccgtt 1080	)
taaccgaggg	agettgteet gagagetaeg gaetteaagt ggeaeteatg getggaatae 1140	)
caaaccaagt	ggttgaaaca gcatcaggtg ctgctcaagc catgaagaga tcaattgggg 1200	)
aaaacttcaa	gtcaagtgag ctaagatetg agtteteaag tetgeatgaa gaetggetea 1260	)
agtcattggt	gggtatttct cgagtcgccc acaacaatgc ccccattggc gaagatgact 1320	)
acgacacttt	gtuucgetta tggeatgaga teaaateete ttaurghght eecaaataac 1380	)

ccāāā	13	885
<210>	28	
<211>	34	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	MSH6 specific primer 2S8 for PCR using cDNA of Arabidopsis thaliana ecotype Columbia	
<400>	26	
atcccgggtt a	atttgggaac acagtaagag gatt	34
<210>	29	
<211>	27	
<212>	DNA .	
<213>	Artificial sequence	
<220>		
<223>	MSH6 specific primer S82 for PCR using cDNA of Arabidopsis thaliana ecotype Columbia	
<400>	29	
gcgttcgatc	atcagectet gtgttge	27
<210>	30	
<211>	3606	
<212>	DNA	
<213>	Arabidopsis thaliana ecotype Columbia	
<220>		
<221>	CDS	
<222>	(142)(3468)	
<223>	AtMSH6 full-length cDNA and deduced sequence of the encode polypeptide	ed.
<400>	30	
aaaagttgag	ccctgaggag tatcgtttcc gccatttcta cgacgcaagg cgaaaatttt	60
tggcgccaat	ctttccccc tttcgaattc tctcagctca aaacatcgtt tctctctcac 12	20
tctctctcac	aattecaaaa a atg cag cgc cag aga teg att ttg tet tte  Met Gln Arg Gln Arg Ser Ile Leu Ser Phe  1 5 10	71

ttc Phe	caa Gln	aaa Lys	ccc Pro	acc Thr 15	gcg Ala	gcg Ala	act Thr	acg Thr	aag Lys 20	ggt Gly	ttg Leu	gtt Val	tcc Ser	ggc Gly 25	gat Asp		219
								gga Gly 35									267
gaa Glu	Gly	gat Asp 45	gct Ala	aaa Lys	ggc Gly	gac Asp	gct Ala 50	tct Ser	gta Val	cgt Arg	ttt Phe	gct Ala 55	gtt Val	tcg Ser	aaa Lys		315
								gat Asp									363
cgt Arg 75	cgt Arg	gtc Val	ctg Leu	ccg Pro	tct Ser 80	gga Gly	ttt Phe	aag Lys	ccg Pro	gct Ala 85	gaa Glu	tcc Ser	gcc Ala	gst Gly	gat Asp 90		411
gct Ala	tcg Ser	tcc Ser	ctg Leu	ttc Phe 95	tcc Ser	aat Asn	att Ile	atg Met	cat His 100	aag Lys	ttt Phe	gta Val	aaa Lys	gtc Val 105	gat Asp	<b>;</b> .	459
gat Asp	cga Arg	gat Asp	tgt Cys 110	tct Ser	gga Gly	gag Glu	agg Arg	agc Ser 115	cga Arg	gaa Glu	gat Asp	gtt Val	gtt Val 120	ccg Pro	ctg Leu		507
								gct Ala									555
cgt Arg	tcc Ser 140	aat Asn	aat Asn	ggt Gly	aaa Lys	act Thr 145	caa Gln	gaa Glu	aga Arg	aac Asn	cat His 150	gct Ala	ttt Phe	agt Ser	ttc Phe		603
	Gly					Arg		gta Val			Ile				ggc Gly 170		651
					Glu			ggg Gly		Arg							699
				Leu					Thr					Lys	gtt Val		747
			Asp					, Leu					Asp		gtt Val		795

tgt Cys	gga Gly 220	gag Glu	aag Lys	aaa Lys	gaa Glu	gta Val 225	aac Asn	gaa Glu	gga Gly	acc Thr	aaa Lys 230	ttt Phe	gaa Glu	tgg Trp	ctt Leu		843
gag Glu 235	tct Ser	tct Ser	cga Arg	atc Ile	agg Arg 240	gat <b>Asp</b>	gcc Ala	aat Asn	aga Arg	aga Arg 245	cgt Arg	cct Pro	gat Asp	gat Asp	ecc Pro 250		891
ctt Leu	tac Tyr	gat Asp	aga Arg	aag Lys 255	acc Thr	tta Leu	cac His	ata Ile	cca Pro 260	cct Pro	gat Asp	gtt Val	ttc Phe	aag Lys 265	aaa Lys		939
			tca Ser 270													_	987
gac Asp	att Ile	gtg Val 285	ctt Leu	ttc Phe	ttt Phe	aaa Lys	gtg Val 290	Gly 999	aaa Lys	ttt Phe	tat Tyr	gag Glu 295	ctg Leu	tat Tyr	gag Glu		1035
cta Leu	gat Asp 300	gcg Ala	gaa Glu	tta Leu	ggt Gly	cac His 305	aag Lys	gag Glu	ctt Leu	gac Asp	tgg Trp 310	aag Lys	atg Met	acc Thr	atg Met		1083
agt Ser 315	ggt Gly	gtg Val	gga Gly	aaa Lys	tgc Cys 320	aga Arg	cag Gln	gtt Val	ggt Gly	atc Ile 325	tct Ser	gaa Glu	agt Ser	GJÅ āāā	ata Ile 330		1131
gat Asp	gag Glu	gca Ala	gtg Val	caa Gln 335	aag Lys	cta Leu	tta Leu	gct Ala	cgt Arg 340	gga Gly	tat Tyr	aaa Lys	gtt Val	gga Gly 345	cga Arg		1179
atc Ile	gag Glu	cag Gln	cta Leu 350	Glu	aca Thr	tct Ser	gac Asp	caa Gln 355	gca Ala	aaa Lys	gcc Ala	aga Arg	ggt Gly 360	gct Ala	aat Asn		1227
act Thr	ata Ile	att Ile 365	Pro	agg Arg	aag Lys	cta Leu	gtt Val 370	Gln	gta Val	tta Leu	act Thr	cca Pro 375	Ser	aca Thr	gca Ala		1275
ago Ser	gag Glu 380	Gly	aac Asn	ato Ile	ggg Gly	cct Pro 385	Asp	gcc Ala	gtc Val	cat His	Leu 390	Leu	gct Ala	ata Ile	aaa Lys		1323
gag Glu 395	Ile	aaa Lys	atg Met	g gag	t Leu 400	Glr	aag Lys	tgt Cys	tca Ser	act Thr	Val	tat Tyr	gga Gly	ttt Phe	gct Ala 410		1371
t t t Phe	gtt Val	gad As <u>r</u>	tgt Cys	gct Ala	a Ala	ttg Leu	agg Arg	j ttt j Phe	tgg Trp 420	val	Gly	tcc Ser	atc : Ile	ago Ser 425	gat Asp		1419

gat Asp	gca Ala	tca Ser	tgt Cys 430	gct Ala	gct Ala	ctt Leu	gga Gly	gcg Ala 435	tta Leu	t tg Leu	atg Met	cag Gln	gtt Val 440	tct Ser	cca Pro	1467
aag Lys	gaa Glu	gtg Val 445	tta Leu	tat Tyr	gac Asp	agt Ser	aaa Lys 450	Gly 999	cta Leu	tca Ser	aga Arg	gaa Glu 455	gca Ala	caa Gln	aag Lys	1515
gct Ala	cta Leu 460	agg Arg	aaa Lys	tat Tyr	acg Thr	ttg Leu 465	aca Thr	Gly ggg	tct Ser	acg Thr	gcg Ala 470	gta Val	cag Gln	ttg Leu	gct Ala	1563
cca Pro 475	gta Val	cca Pro	caa Gln	gta Val	atg Met 480	ggg Gly	gat Asp	aca Thr	gat Asp	gct Ala 485	gct Ala	gga Gly	gtt Val	aga Arg	aat Asn 490	1611
ata Ile	ata Ile	gaa Glu	tct Ser	aac Asn 495	gga Gly	tac Tyr	ttt Phe	aaa Lys	ggt 500	t <i>c</i> t Ser	tct Ser	gaa Glu	tca Ser	tgg Trp 505	aac Asn	1659
tgt Cys	gct Ala	gtt Val	gat Asp 510	ggt Gly	cta Leu	aat Asn	gaa Glu	tgt Cys 515	gat Asp	gtt Val	gcc Ala	ctt Leu	agt Ser 520	gct Ala	ctt Leu	1707
gga Gly	gag Glu	cta Leu 525	Ile	aat Asn	cat His	ctg Leu	ser 530	agg Arg	cta Leu	aag Lys	cta Leu	gaa Glu 535	gat Asp	gta Val	ctt Leu	1755
aag Lys	cat His 540	Gly	gat Asp	att Ile	ttt Phe	cca Pro <b>545</b>	tac Tyr	caa Gln	gtt Vai	tac Tyr	agg Arg 550	Gly	tgt Cys	ctc Leu	aga Arg	1803
att Ile 555	: Asp	ggc Gly	cag Glr	acg Thr	atg Met 560	Val	aat Asn	ctt Leu	gag Glu	ata Ile 565	Phe	aac Asn	aat Asn	agc Ser	tgt Cys 570	1851
gat Asp	ggt Gly	ggt Gly	cct Pro	tca Ser 575	Gly	acc Thr	ttg Lev	tac Tyr	aaa Lys 580	Tyr	ctt Lev	gat Asp	aac Asn	tgt Cys 585	Val	1899
agi Se:	r Pro	a act	ggt r Gly	y Lys	g cga	a cto	tta Lev	a agg a Arg 595	Ası	tgg Trp	g ato	tgo Cys	cat His	Pro	ctc Leu	1947
aa Ly	a ga s As	t gt p Va 60	1 G1	a ago u Se	c ato	c aat e Asr	aaa Lys	s Arg	g Cti	t gat u Ası	gta p Vai	a gt: 1 Va: 61:	l Glu	a gaa 1 Glu	a ttc ı Phe	1995
ac Th	g gc r Al 62	a As	c tc n Se	a ga r Gl	a ag u Se	t ato	t Gl	a ato n Ile	ac e Th	t gg r Gl	c ca y Gl: 63	n Ty	t cto	c cad u Hi:	c aaa s Lys	2043

ctt Leu		-		-	aga Arg	-		_	-				_	_	-	2091
635					640					645					650	
		-			ttg Leu		-		_					_		2139
	_	_		•	ttt Phe								-	_		2187
	_	_	_	_	gct Ala		_	_	_			_	_	_	_	2235
				_						_			_		cta Leu	2283
-					caa Gln 720		-	-	-		_	_	_			2331
		_			gat Asp			-	-		_	-				2379
			-		ttt Phe		-	_	-						_	2427
				_	tgc Cys		-	-	_	-			_		_	2475
_	_			_	gga Gly	_	-	-			_				_	2523
					cag Gln 800											2571
					cat											2619
					ata Ile											2667

														ggc Gly			2715
					-	-		_	_	_	_			gcc Ala			2763
		-			-				_	-				gtg Val	-		2811
							-		-	_		_		gga Gly 905			2859
-			_	_	_	_						_		cag Gln			2907
_		_	_						_	_	_			gga Gly		· (c.	2955
_			_							_				cac His			3003
_			_		_		_			_				cac His			3051
		_	•						_	•		_		cac His 985	_		3099
-		_						_				Arg		tgt C <b>ys</b>	-		3147
	Asp					Tyr					Gly	_	_	cct Pro			3195
Ser					Val			-	_	Gly				caa Gln	-		3243
	Glu			Ser		Ala			Ala	_	_	_		att Ile			3291

			Lys					Arg			ttc Phe		Ser			3339
_	-	Trp		_		_	Val				cga Arg	Val	_			3387
	Ala				_	Asp	_		_		ttg Leu 1		-			3435
His					Ser		tgt Cys	_			taaa	tggo	ta			3478
tgac	ataa	.ca c	tato	tgaa	g ct	.cgtt	aagt	ctt	ttgo	ctc	tctg	atgt	tt a	ttcc	tctta	3538
aaaa	atgo	tt a	tata	tcaa	a aa	attg	ittt	cto	gatt	aaa	aaaa	aaaa	iaa a	aaaa	aaaaa ,3	3598
aaaa	aaaa	L						•							, š	3606
<210 <211 <212 <213 <223	l> !> !>			abido	psis cide			na ec	oty <u>r</u>	e Co	olumb	oia				
<400	)>		31													
Met 1	Gln	Arg	Gln	Arg 5	Ser	Ile	Leu	Ser	Phe 10	Phe	Gln	Lys	Pro	Thr 15	Ala	
Ala	Thr	Thr	Lys 20	Gly	Leu	Val	Ser	Gly 25	Asp	Ala	Ala	Ser	Gly 30	Gly	Gly	
Gly	Ser	Gly 35	Gly	Pro	Arg	Phe	Asn 40	Val	Arg	Glu	Gly	Asp 45	Ala	Lys	Gly	
Asp	Ala 50	Ser	Val	Arg	Phe	Ala 55	Val	Ser	Lys	Ser	Val 60	Asp	Glu	Val	Arg	
Gly 65	Thr	Asp	Thr	Pro	Pro 70	Glu	Lys	Val	Pro	Arg 75	Arg	Val	Leu	Pro	Ser 80	
Gly	Phe	Lys	Pro	Ala 85	Glu	Ser	Ala	Gly	Asp 90	Ala	Ser	Ser	Leu	Phe 95	Ser	
Asn	Ile	Met	His 100	Lys	Phe	Val	Lys	Val 105	Asp	Asp	Arg	Asp	Cys 110	Ser	Gly	
Glu	Arg	Ser	Arg	Glu	Asp	Val	Val 120		Leu	Asn	Asp	ser 125	Ser	Leu	Cys	

WO 99/19492 PCT/EP98/06977

Met Lys Ala Asn Asp Val Ile Pro Gln Phe Arg Ser Asn Asn Gly Lys Thr Gln Glu Arg Asn His Ala Phe Ser Phe Ser Gly Arg Ala Glu Leu Arg Ser Val Glu Asp Ile Gly Val Asp Gly Asp Val Pro Gly Pro Glu Thr Pro Gly Met Arg Pro Arg Ala Ser Arg Leu Lys Arg Val Leu Glu Asp Glu Met Thr Phe Lys Glu Asp Lys Val Pro Val Leu Asp Ser Asn Lys Arg Leu Lys Met Leu Gln Asp Pro Val Cys Gly Glu Lys Lys Glu. . Val Asn Glu Gly Thr Lys Phe Glu Trp Leu Glu Ser Ser Arg Ile Arg Aso Ala Asn Arg Arg Pro Asp Asp Pro Leu Tyr Asp Arg Lys Thr Leu His Ile Pro Pro Asp Val Phe Lys Lys Met Ser Ala Ser Gln Lys Gln Tyr Trp Ser Val Lys Ser Glu Tyr Met Asp Ile Val Leu Phe Phe Lys Val Gly Lys Phe Tyr Glu Leu Tyr Glu Leu Asp Ala Glu Leu Gly His Lys Glu Leu Asp Trp Lys Met Thr Met Ser Gly Val Gly Lys Cys Arg Gln Val Gly Ile Ser Glu Ser Gly Ile Asp Glu Ala Val Gln Lys Leu Leu Ala Arg Gly Tyr Lys Val Gly Arg Ile Glu Gln Leu Glu Thr Ser Asp Gln Ala Lys Ala Arg Gly Ala Asn Thr Ile Ile Pro Arg Lys Leu Val Gln Val Leu Thr Pro Ser Thr Ala Ser Glu Gly Asn Ile Gly Pro Asp Ala Val His Leu Leu Ala Ile Lys Glu Ile Lys Met Glu Leu Gln Lys Cys Ser Thr Val Tyr Gly Phe Ala Phe Val Asp Cys Ala Ala 

Leu Arg Phe Trp Val Gly Ser Ile Ser Asp Asp Ala Ser Cys Ala Ala 420 Leu Gly Ala Leu Leu Met Gln Val Ser Pro Lys Glu Val Leu Tyr Asp Ser Lys Gly Leu Ser Arg Glu Ala Gln Lys Ala Leu Arg Lys Tyr Thr 455 Leu Thr Gly Ser Thr Ala Val Gln Leu Ala Pro Val Pro Gln Val Met 470 475 Gly Asp Thr Asp Ala Ala Gly Val Arg Asn Ile Ile Glu Ser Asn Gly 485 490 Tyr Phe Lys Gly Ser Ser Glu Ser Trp Asn Cys Ala Val Asp Gly Leu 500 505 Asn Glu Cys Asp Val Ala Leu Ser Ala Leu Gly Glu Leu Ile Asn His 520 Leu Ser Arg Leu Lys Leu Glu Asp Val Leu Lys His Gly Asp Ile Phe 530 535 Pro Tyr Gln Val Tyr Arg Gly Cys Leu Arg Ile Asp Gly Gln Thr Met 545 550 Val Asn Leu Glu Ile Phe Asn Asn Ser Cys Asp Gly Gly Pro Ser Gly 570 Thr Leu Tyr Lys Tyr Leu Asp Asn Cys Val Ser Pro Thr Gly Lys Arg 585 Leu Leu Arg Asn Trp Ile Cys His Pro Leu Lys Asp Val Glu Ser Ile 595 600 Asn Lys Arg Leu Asp Val Val Glu Glu Phe Thr Ala Asn Ser Glu Ser 615 Met Gln Ile Thr Gly Gln Tyr Leu His Lys Leu Pro Asp Leu Glu Arg 630 Leu Leu Gly Arg Ile Lys Ser Ser Val Arg Ser Ser Ala Ser Val Leu 645 650 Pro Ala Leu Leu Gly Lys Lys Val Leu Lys Gln Arg Val Lys Ala Phe 660 665 Gly Gln Ile Val Lys Gly Phe Arg Ser Gly Ile Asp Leu Leu Leu Ala 685

Leu din Lys Glu Ser Asn Met Met Ser Leu Leu Tyr Lys Leu Cys Lys

700

695

Leu 705	Pro	Ile	Leu	Val	Gly 710	Lys	Ser	Gly	Leu	Glu 715	Leu	Phe	Lau	Ser	Gln 720
Phe	Glu	Ala	Ala	Ile 725	Asp	Ser	Asp	Phe	Pro 730	Asn	Tyr	Gln	Asn	Gln 735	Asp
Val	Thr	Asp	Glu 740	Asn	Ala	Glu	Thr	Leu 745	Thr	Ile	Leu	Ile	Glu 750	Leu	Phe
Ile	Glu	Arg 755	Ala	Thr	Gln	Trp	Ser 760	Glu	Val	Ile	His	Thr 765	Ile	Ser	Cys
Leu	<b>Asp</b> 770	Val	Leu	Arg	Ser	Phe 775	Ala	Ile	Ala	Ala	Ser 780	Leu	Ser	Ala	Gly
Ser 785	Met	Ala	Arg	Pro	Val 790	Ile	Phe	Pro	Glu	Ser 795	Glu	Ala	Thr	Asp	Gln 800
Asn	Gln	Lys	Thr	Lys 805	Gly	Pro	Ile	Leu	Lys 810	Ile	Gln	Gly	Leu	Trp 815	His
Pro	Phe	Ala	Val 820	Ala	Ala	qsA	Gly	Gln 825	Leu	Pro	Val	Pro	Asn 830	Asp	Ile
Leu	Leu	Gly 835	Glu	Ala	Arg	Arg	Ser 840	Ser	Gly	Ser	Ile	His 845	Pro	Arg	Ser
Leu	Leu 850	Leu	Thr	Gly	Pro	Asn 855	Met	Gly	Gly	Lys	Ser 860	Thr	Leu	Leu	Arg
Ala 865	Thr	Cys	Leu	Ala	Val 870	Ile	Phe	Ala	Gln	Leu 875	Gly	Cys	Tyr	Val	Pro 880
Cys	Glu	Ser	Cys	Glu 885	Ile	Ser	Leu	Val	Asp 890	Thr	Ile	Phe	Thr	Arg 895	Leu
Gly	Ala	Ser	Asp 900	Arg	Ile	Met	Thr	Gly 905	Glu	Ser	Thr	Phe	Leu 910	Val	Glu
Cys	Thr	Glu 915	Thr	Ala	Ser	Val	Leu 920	Gln	Asn	Ala	Thr	Gln 925	Asp	Ser	Leu
Val	Ile 930	Leu	Asp	Glu	Leu	Gly 935	Arg	Gly	Thr	Ser	Thr 940	Phe	Asp	Gly	Tyr
Ala 945	Ile	Ala	Tyr	Ser	Val 950	Phe	Arg	His	Leu	Val 955	Glu	Lys	Val	Gln	Cys 960
Arg	Met	Leu	Phe	Ala 965	Thr	His	Tyr	His	Pro 970	Leu	Thr	Lys	Glu	Phe 975	Ala
Ser	His	Pro	Arg 980	Val	Thr	Ser	Lys	His 985	Met	Ala	Cys	Ala	Phe 990	Lys	Ser

PCT/EP98/06977

Arg Ser	Asp 995	Tyr Gln	Pro Arg	Gly Cy 1000	rs Asp	Gin As	1005		Fife	Leu	
Tyr Arg 1010	Leu	Thr Glu	Gly Ala 1015	Cys Pr	o Glu	Ser Ty		Leu	Gln	Val	
Ala Leu 1025	Met		Ile Pro 1030	Asn Gl		Val G1 1035	u Thr	Ala		Gly 040	
Ala Ala	Gln	Ala Met 1045	Lys Arg	Ser Il	e Gly 1050		n Phe		Ser 055	Ser	
Glu Leu	_	Ser Glu .060	Phe Ser	Ser Le		Glu As		Leu 1070	Lys	Ser	
	Gly 1075	Ile Ser	Arg Val	Ala Hi 1080	.s Asn	Asn Al	a Pro 1085		Gly	Glu	
Asp Asp 1090		Asp Thr	Leu Phe 1095		eu Trp	His G		Lys	Ser	Ser	÷.
Tyr Cys	Val	Pro Lys									
<210>		32									
<211>		24									
<212>		DNA	-i-l								
<213>		Artill	cial seq	uence						•	
<220>											
<223>			d primer atellite		CR amp	olifica	tion o	f ATH	(GEN)	EΑ	
<400>		32									
accatgo	ata	gcttaaac	ett cttg								24
<210>		33									
<211>		22									
<212>		DNA									
<213>		Artifi	cial sec	quence							
<220>		•									
<223>			se prime: satellite		CR am	plifica	tion o	of AT	HGEN	EA	
<400>		33									
acataa	ccac	aaatagg	ggt gc								22

29

gototagatt totogotota agactotot

<210>	34
<211>	18
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Forward primer DMCIN-A for PCR on genomic DNA of Arabidopsis
	thaliana ssp. Landsberg erecta "Ler"
<400>	34
gaagcgatat t	gttcgtg 18
<210>	35
<211>	18
<212>	DNA
<213>	Artificial sequence
<220>	•
<223>	Reverse primer DMCIN-B for PCR on genomic DNA of Arabidopsis
	thaliana ssp. Landsberg erecta "Ler"
<400>	35
<b>aga</b> ttgcgag a	acattcc 18
<210>	36
<211>	31
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Forward primer DMCIN-1 for PCR on genomic DNA of Arabidopsis
	thaliana ssp. Landsberg erecta "Ler"
<400>	36
acgcgtcgac t	cagetatga gattactegt g 31
<210>	17
	37
<211>	29
<212>	
<213>	Artificial sequence
222	
<220>	
<223>	Thereads making DMCTTM I have DCD on managin DMT of Aughidensia
	Reverse primer DMCIN-2 for PCR on genomic DNA of Arabidopsis
	thaliana ssp. Landsberg erecta "Ler"
	thaliana ssp. Landsberg erecta "Ler"
<400>	

<210>	38	
<211>	32	
<212>	DNA	
<213>	Artificial sequence	
<220>	,	
<223>	Forward primer DMCIN-3 for PCR on genomic DNA of Arabidopsisthaliana ssp. Landsberg erecta "Ler"	3
<400>	38	
gctctagagc	ttctcttaag taagtgattg at 3.	2
<210>	39	
<211>	48	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer DMCIN-4 for PCR on genomic DNA of Arabidopsichaliana ssp. Landsberg erecta "Ler"	5
<400>	39	
tcccccgggc	tcgagagatc tccatggttt cttcagctct atgaatcc 4	8
<210>	40	
<211>	26	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer DMCla for PCR on genomic DNA of Arabidopsis thaliana ssp. Landsberg erecta "Ler"	
<400>	40	
acgcgtcgac	gaattcgcaa gtgggg 2	6
<210>	41	
<211>	38	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer DMClb for PCR on genomic DNA of Arabidopsis thaliana ssp. Landsberg erecta "Ler"	
<400×		

recategaga	tetecegggt accgatttge ttegaggg	38
<210>	42	
<211>	20	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of ATEAT1 SSLP marker	in
	Arabidopsis thaliana subspecies	
<400>	42	
gccactgcgt	gaatgatatg	20
<210>	43	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
<220>	, ·	
<223>	Reverse primer for PCR amplification of ATEAT1 SSLP marker	in
	Arabidopsis thaliana subspecies	
<400>	43	
cgaacagcca	acattaattc cc	22
<210>	44	
<211>	18	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA63 SSLP marker	in
	Arabidopsis thaliana subspecies	
<400>	44	
aaccaaggca	cagaageg	18
<210>	45	
<211>	18	
<212>	DNA	
<213>	Artificial sequence	
	-	
<220>		
<223>	Reverse primer for PCR amplification of NGA63 SSLP marker Arabidopsis thaliana subspecies	in

<400>	45	
acccaagtga	tegecace	18
	16	
<210>	46	
<211>	21	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA248 SSI Arabidopsis thaliana subspecies	lP marker in
<400>	46	
taccgaacca	aaacacaaag g	21
		•
<210>	47	÷.
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of NGA248 SS Arabidopsis thaliana subspecies	LP marker ir
<400>	47	
tctgtatctc	ggtgaattet ee	22
<210>	48	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA128 SS	LP marker i
62237	Arabidopsis thaliana subspecies	
<400>	48	
ggtctgttg	a tgtcgtaagt cg	22
<210>	49	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
<220>		

<223>	Reverse primer for PCR amplification of Arabidopsis thaliana subspecies	NGA128	SSLP	marker in
<400>	49			
atcttgaaac c	tttagggag gg			22
<210>	50			
<211>	22			
<212>	DNA			
<213>	Artificial sequence			
<220>				
<223>	Forward primer for PCR amplification of Arabidopsis thaliana subspecies	NGA280	SSLP	marker in
<400>	50			
ctgatctcac g	gacaatagt gc			22
	·	•		
<210>	51	•		
<211>	20			
<212>	DNA	•		
<213>	Artificial sequence			
	-			
<220>	,			
<223>	Reverse primer for PCR amplification of	NGA280	SSLP	marker in
	Arabidopsis thaliana subspecies			
<400>	51			
<400>	51			
ggctccataa a	aagtgcacc			20
<210>	52			
<211>	21			
<212>	DNA			
<213>	Artificial sequence			
<220>				
<223>	Forward primer for PCR amplification of	NGA111	SSLP	marker in
	Arabidopsis thaliana subspecies			
<400>	52			
ctccagttgg a	aagctaaagg g			21
<210>	53			
<211>	21			
<212>	UNA			
<213>	Artificial sequence			

PCT/EP98/06977

<220>		
<223>	Reverse primer for PCR amplification of NGAlll SSLP marke Arabidopsis thaliana subspecies	r i
<400>	53	
tgttttttag	gacaaatggc g	21
<210>	54	
<211>	20	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA168 SSLP market Arabidopsis thaliana subspecies	r in
<400>	54	
ccttcacatc	caaaacccac	20
•		
<210>	55	
<211>	20	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of NGA168 SSLP marker Arabidopsis thaliana subspecies	in
<400>	55	
gcacataccc	acaaccagaa	20
<210>	56	
<211>	20	
<212>	DNA	
<213>		
(213)	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA1126 SSLP marks in Arabidopsis thaliana subspecies	er
<400>	56	
cgctacgctt	ttcggtaaag	20

<210>	57
<211>	20
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Reverse primer for PCR amplification of NGA1126 SSLP marker
	in Arabidopsis thaliana subspecies
<400>	57
gcacagtcca	agtcacaacc 20
<210>	58
<211>	20
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Forward primer for PCR amplification of NGA361 SSLP marker in Arabidopsis thaliana subspecies
<400>	58
<b>aaaga</b> gatga	gaatttggac 20
<210>	 59
<210 <i>&gt;</i>	23
<211>	DNA
<213>	Artificial sequence
<220>	•
<223>	Reverse primer for PCR amplification of NGA361 SSLP marker in Arabidopsis thaliana subspecies
<400>	59
acatatcaat	atattaaagt agc 23
<210>	60
<211>	18
<212>	DNA
<213>	Artificial sequence
~~137	Wrettrerar seduction
<220>	
<223>	Forward primer for PCR amplification of NGA168 SSLP marker in Arabidopsis thaliana subspecies
<400>	60

40

tegtetactg	cactgccg	18
<210>	61	
<211>	22	
<212>	DNA	
<212>		
<213>	Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of NGA168 SSLP marker Arabidopsis thaliana subspecies	r in
<400>	61	
gaggacatgt	ataggageet eg	22
<210>	62	
<211>	20	
<211><212>	DVA	
<212>	Artificial sequence	
(213)	Altilitial Sequence	
<220>		
<223>	Forward primer for PCR amplification of AthBIO2 SSLP marker in Arabidopsis thaliana subspecies	er
<400>	62 	
tgacctcctc	ttccatggag	20
<210>	63	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of AthBIO2 SSLP marks	er
	in Arabidopsis thaliana subspecies	
<400>	63	
ttaacagaaa	cccaaagctt tc	22
<210>	64	
<211>	21	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Powered arimon for DCD amplification of Athenrous corp	~ - داء
~6637	Forward primer for "CR amplification of AthUBIQUE SSLP may in Arabidopsis thaliana subspecies	KET

<400>	64				
aggcaaatgt	ccatttcatt g				21
	•				
<210>	65				
<211>	20				
<212>	DNA				
<213>	Artificial sequence				
	•				
<220>					
<223>	Reverse primer for PCR amplification	of	AthUBIQ	UE SSLP	marker
	in Arabidopsis thaliana subspecies				
<400>	65				
acgacatggc	agatttctcc				20
			•		
210					
<210><211>	66		•	ές.	
<211>	21			•	
	DNA				
<213>	Artificial sequence				
<220>					
<223>	Forward primer for PCR amplification	of	NGA172	SSLP ma	rker in
	Arabidopsis thaliana subspecies	-			
<400>	66				
agctgcttcc	ttatagcgtc c				21
-210-	67				
<210> <211>	67 19				
<211>	DNA				
<213>	Artificial sequence				
(213)	Altilitial Sequence				
<220>					
<223>	Reverse primer for PCR amplification	of	NGA172	SSLP ma	rker in
	Arabidopsis thaliana subspecies	-		J001	2,102 21
<400>	67				
catccgaatg	ccattgttc				19
٠.					
<210>	68				
<211>	21				
<212>	DNA				
<213>	Artificial sequence				
<220>					

<223>	Forward primer for PCR amplification Arabidopsis thaliana subspecies	of	NGA126	SSLP	marker	in
<400>	63					
gaaaaaacgc	tactttcgtg g					21
<210>	69					
<211>	22					
<212>	DNA					
<213>	Artificial sequence					
<220>						
<223>	Reverse primer for PCR amplification ( Arabidopsis thaliana subspecies	of	NGA126	SSLP	marker	ın
<400>	69				•	
caagagcaat	atcaagagca gc					22
				. '5		
<210>	70					
<211><212>	20 DNA					
<212>	Artificial sequence					
(213)	Arciriciar sequence					
<220>						_
<223>	Forward primer for PCR amplification Arabidopsis thaliana subspecies	of	NGA162	SSLP	marker	in
<400>	70					
catgcaattt	gcatctgagg					20
<210>	71					
<211>	22					
<212>	DNA					
<213>	Artificial sequence					
<220>						
<223>	Reverse primer for PCR amplification Arabidopsis thaliana subspecies	of	NGA162	SSLP	marker	in
<400>	71					•
ctctgtcact	cttttcctct gg					22
.210-	7.2					
<210>	72					
<211>	21 DNA					
<212><213>	DNA Artificial sequence					
42137	wrettrerat sedaeuce					

<220>		
<223>	Forward primer for PCR amplification of NGA6 SSLP marker : Arabidopsis thaliana subspecies	in
<400>	72	
tggatttctt	cctctcttca c	21
<210>	73	
<211>	21	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of NGA6 SSLP marker in Arabidopsis thaliana subspecies	in
<400>	73	
atggagaagc	ttacactgat c	21
<210>	74	
<211>	20	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA12 SSLP marker Arabidopsis thaliana subspecies	in
<400>	74	
aatgttgtcc	tcccctc	20
-210-	35	
<210><211>	75 22	
<211>	DNA	
<213>	Artificial sequence	
<220>	Develope puises for non-amplification of Monte con-	
<223>	Reverse primer for PCR amplification of NGA12 SSLP marker Arabidopsis thaliana subspecies	ın
<400>	75	
tgatgctctc	tgaaacaaga gc	22

<210>	76
<211>	21
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Forward primer for PCR amplification of NGA8 SSLP marker in Arabidopsis thaliana subspecies
<400>	76
gagggcaaat	ctttatttcg g
<210>	77
<211>	22
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Reverse primer for PCR amplification of NGA8 SSLP marker in Arabidopsis thaliana subspecies
<400>	77
tggctttcgt	ttataaacat cc 22
<210>	70
<210 <i>&gt;</i>	78
<211> <212>	21
	DNA
<213>	Artificial sequence
<220>	
<223>	Forward primer for PCR amplification of NGA1107 SSLP marker in Arabidopsis thaliana subspecies
<400>	78
gcgaaaaaac	aaaaaaatcc a 21
-210.	70
<210>	79
<211>	21
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Reverse primer for PCR amplification of NGA1107 SSLP marker in Arabidopsis thaliana subspecies
<400>	79

cgacgaatcg	acagaattag g	21
<210>	80	
<211>	21	
<212>	DNA	
<213>	Artificial sequence	
10207	viciliciai seducuce	
<220>		
<223>	Forward primer for PCR amplification of NGA225 SSLP marker Arabidopsis thaliana subspecies	in
<400>	80	
gaaatccaaa	tcccagagag g	1
<210>	81	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Payarsa primar for pap	
	Reverse primer for PCR amplification of NGA225 SSLP marker : Arabidopsis thaliana subspecies	in
<400>	81	
tctccccact a	agttttgtgt cc 2	2
<210>	82	
<211>	19	
<212>	DNA	
<213>	Artificial sequence	
<220> <223>		
(223)	Forward primer for PCR amplification of NGA249 SSLP marker i Arabidopsis thaliana subspecies	.n
<400>	82	
taccgtcaat t	icategee	<b>,</b>
<210>	83	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
-220-		
<220> <223>	Description of the second of t	
~ & & J 7	Reverse primer for PCR amplification of NGA249 SSLP marker i Ambidopsis thaliana subspecies	n

	· · ·
<400>	83
ggatecetaa	ctgtaaaatc cc 22
<210>	84
<211>	22
<212>	DNA .
<213>	Artificial sequence
	*
<220>	
<223>	Forward primer for PCR amplification of CA72 SSLP marker in Arabidopsis thaliana subspecies
<400>	84
aaccccagta a	ccaaacaca ca 22
<210>	85
<211>	20
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Reverse primer for PCR amplification of CA72 SSLP marker in
	Arabidopsis thaliana subspecies
<400>	85
(100)	
cccagtctaa c	cacgaccac 20
-	
<210>	86
<211>	20
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Forward primer for PCR amplification of NGA151 SSLP marker in
(223)	Arabidopsis thaliana subspecies
	abraoporo charrana sanspectes
<400>	86
gttttgggaa g	ttttgctgg 20
• ,	
<210>	87
<211>	24
<212>	DNA
<213>	Artificial sequence
<220>	
-4637	

<223>	Reverse primer for PCR amplification of NGA151 SSLP marker in Arabidopsis thaliana subspecies
<400>	87
cagtctaaaa	gcgagagtat gatg 24
<210>	88
<211>	22
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Forward primer for PCR amplification of NGA106 SSLP marker in Arabidopsis thaliana subspecies
<400>	88
gttatggagt	ttctagggca cg 22
3 33 5	
<210>	89
<211>	20
<212>	DNA
<213>	Artificial sequence
(213)	ALCITICIAL Sequence
<220>	·
<223>	Reverse primer for PCR amplification of NGA106 SSLP marker in
	Arabidopsis thaliana subspecies
<400>	89
<400>	67
tgccccattt	tgttcttctc 20
<210>	90
<211>	20
<211> <212>	
	DNA
<213>	Artificial sequence
<220>	
<223>	Forward primer for PCR amplification of NGA139 SSLP marker in
	Arabidopsis thaliana subspecies
<400>	90
agagetacea	gatccgatgg 20
<210>	91
<211>	21
<212>	DNA
<213>	Artificial sequence

PCT/EP98/06977

<220>		
<223>	Reverse primer for PCR amplification of NGA139 SSLP marker Arabidopsis thaliana subspecies	i
<400>	91	
ggtttcgttt	cactatccag g	21
<210>	92	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
(213)	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA76 SSLP marker Arabidopsis thaliana subspecies	in
<400>	92	
ggagaaaatg	tcactctcca cc	22
<210>	93	
<211>	20	
<212>	DNA .	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of NGA76 SSLP marker Arabidopsis thaliana subspecies	in
<400>	93	
aggcatggga	gacatttacg	20
<210>	94	
<211>	20	
<212>	D <b>NA</b>	
<213>	Artificial sequence	
	natalata segucite	
<220>		
<223>	Forward primer for PCR amplification of ATHSO191 SSLP mark in Arabidopsis thaliana subspecies	er
<400>	94	
ctccaccaat	catgcaaatg	20

WO 99/19492 PCT/EP98/06977

49

<210>	95	
<211>	21	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of ATHSO191 SSLP marker in Arabidopsis thaliana subspecies	er
<400>	95	
tgatgttgat (	ggagatggtc a	21
<210>	96	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA129 SSLP marker Arabidopsis thaliana subspecies	ir
<400>	96	
tcaggaggaa d	ctaaagtgag gg	22
<210>	97	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of NGA129 SSLP marker Arabidopsis thaliana subspecies	ir
<400>	97	
cacactgaag a	atggtcttga gg	22
<210>	98	
<211>	8062	
<212>	DNA	
<213>	Arabidopsis thaliana ecotype Columbia	
<220>		
<223>	Genomic DNA sequence of AtMSH6	
<400>	97	

ttttttggtt gctaacaata aaggtatacg gttttatgtc atcaatataa ctatatata

aagaaatgaa	agatatatat	tgttttttca	tttatcaaac	aaaacaacaa	gactttttt	120
ttacttttta	cattggtcaa	caaaatacaa	gataaacgac	atogtttaat	catttcccaa	180
ttttacccct	aagtttaaca	cctagaacct	tctccatctt	cgcaagcaca	gcctgattag	240
gaacagettt	accattctca	tattcctgaa	ctacctgagt	cctctcattg	atctgtttcg	300
ccaaatccgc	ttgtgacatc	ttcttctcca	atctcgcttt	ctgtatcatc	aacctcacct	360
ctgctttcac	acgatccatc	gccgcaggct	ctgtttcttc	ttccagcttc	ttcgtgttaa	420
tcaccggaac	cgccgtagat	ttcccctttt	tgttcgaacc	ggcatcgaat	ttcttaaccg	480
tttgaaccgc	gacaccgttt	ctcagagctg	cgttaaccgc	tttcggatcg	cgtaggtett	540
ggctcttttg	ttttgatttg	tggagaacta	ctggttccca	gtcttgtgtt	actgctcctg	600
ggtatctgct	cggcatcgtc	gatgaattga	gagaaaggaa	caacgcgaaa	attttatta'a	660
tetgagtttt	gaaattgaga	aacgatgaag	atgaagaatg	ttgttgagag	gattgtgata	720
tttatatata	cgaagattgg	tttctggaga	attcgatcat	ctttttctcc	attttcgtct	780
ctggaacgtt	cttagagatg	attgacgacg	tgtcattatc	tgatttgcag	ttaaccaatg	840
ctttttgggt	tggattcgtg	gtacaccata	ttatccgatt	tggctcaatg	gttttatata	900
aatttggttt	teggtteggt	tatgagttat	cattaaaatt	aagctaacca	aaaattttcg	960
taaaatttat	ttcggtttca	attcggatcc	cttacttcca	gaaccgaatt	attcgaaacc	1020
ggggttagcc	gaaccgaata	ccaatgeetg	attgactcgt	tggctagaaa	gatccaacgg	1080
tatacaataa	tagaacataa	atcggacggt	catcaaagcc	tcaaagagtg	aacagtcaac	1140
aaaaaagtt	gagccctgag	gagtatcgtt	tccgccattt	ctacgacgca	aggcgaaaat	1200
ttttggcgcc	aatctttccc	ccctttcgaa	ttctctcagc	tcaaaacatc	gtttctctct	1260
cactetetet	Cacaattcca	aaaaatgcag	cgccagagat	cgattttgtc	tttcttccaa	1320
aaacccacgg	cggcgactac	gaagggtttg	gtttccggcg	atgctgctag	cggcgggggc	1380
ggcagcggag	accacgattt	aatgtgaagg	aaggggatgc	taaaggcgac	gcttctgtac	1440
gttttgctgt	ttcgaaatct	gtcgatgagg	ttagaggaac	ggatactcca	ccggagaagg	1500
ttccgcgtcg	tgtcctgccg	tctggattta	agccggctga	atccgccggt	gatgettegt	1560
ccctgttctc	caatattatg	cataagtttg	taaaagtcga	tgatcgagat	tgttctggag	1620
agaggtacta	atcttcgatt	ctcttaattt	tgttatcttt	agctggaaga	agaagattcg	1680

rotaatttgt	tgtattcgtt	ggagagattc	tgattactgc	attggatcgt	tgtttacaaa	1740
ttttcaggag	ccgagaagat	gttgttccgc	tgaatgattc	atctctatgt	atgaaggcta	1800
atgatgttat	tcctcaattt	cgttccaata	atggtaaaac	tcaagaaaga	aaccatgctt	1860
ttagtttcag	tgggagagct	gaacttagat	cagtagaaga	tataggagta	gatggcgatg	1920
ttcctggtcc	agaaacacca	gggatgcgtc	cacgtgcttc	tcgcttgaag	cgagttctgg	1980
aggatgaaat	gacttttaag	gaggataagg	ttcctgtatt	ggactctaac	aaaaggctga	2040
aaatgctcca	ggatccggtt	tgtggagaga	agaaagaagt	aaacgaagga	accaaatttg	2100
aatggcttga	gtcttctcga	atcagggatg	ccaatagaag	acgtcctgat	gatccccttt	2160
acgatagaaa	gaccttacac	ataccacctg	atgttttcaa	gaaaatgtct	gcatcacaaa	2220
agcaatattg	gagtgttaag	agtgaatata	tggacattgt	gcttttctt	aaagtggtta	2280
gtaactatta	atctagtgtt	caatccattt	cctcaatgtg	atttgttcac	ttacatctgt	2340
ttacgttatg	ctcttctcag	gggaaatttt	atgagctgta	tgagctagat	gcggaattag	2400
gtcacaagga	gcttgactgg	aagatgacca	tgagtggtgt	gggaaaatgc	agacaggtaa	2460
attagttgaa	acaactggcc	tgcttgaatt	attgtgtcta	taaattttga	caccaccttt	2520
tgtttcaggt	tggtatctct	gaaagtggga	tagatgaggc	agtgcaaaag	ctattagctc	2580
gtgggtaagg	gaaccatcat	actttatgga	attcgtttac	tgctacttcg	gctaggattt	2640
aagaaatgga	aatcacttca	agcatcatta	gttaggatcc	tgagaactca	ggatgttttc	2700
ttattcgtta	tataataagt	cttttcatca	aggagtaaca	aacaaaactt	gcacaatatt	2760
tgtgtgctca	ctggcaaggc	atatataccc	agctaacctt	tgctagttca	ctgtagtaac	2820
agttacggat	aatatatgtt	tacttgtatg	tggtaccctc	attttgtctc	tcatggaggc	2880
tttcaagcct	tgtgttgaaa	ctggatagtt	acatatgctt	ccaacagaaa	ctagcatgca	2940
gattcatatg	ctttcctatt	ctactaatta	tgtattgaca	cactcgttgt	ttcttttgaa	3000
agatataaag	ttggacgaat	cgagcagcta	gaaacatctg	accaagcaaa	agccagaggt	3060
gctaatactg	taagttttct	tggataggtc	aaggagagtg	ttgcagactg	tttttgatca	3120
tttcttttc	tgtacattac	tttcatgctg	taattaactc	aatggctatt	ctggtctgat	3180
tatcagataa	ttccaaggaa	gctagttcag	gtattaactc	catcaacagc	aagcgaggga	3240
aacatcgggc	ctgatgccgt	ccatcttctt	gctataaaag	aggtttgtta	tttacttatt	3300

			•			
tatcttatca	tgttcagttc	atccaagtcc	tgaaaaatta	cactettett	taccaatctt	3360
ccatcaagct	gtgtaaagga	tttggaatta	gaaaatcatt	atttgatgct	ttgttttata	3420
tgcaagaggt	tcccttgaaa	agatctgttt	aagattcttt	gcacttgaaa	aattcaatct	3480
ttttaagtga	atcccctact	ttcttacaat	gatcatagtc	tgcaattgca	tgtcaagtaa	3540
tatcattcct	tgttactgca	tcccctctt	tcttaatgac	cattgtctat	gttgtgtttg	3600
tctcgtgtgc	tggagaaaat	gatagctgat	ccaagctgta	cattatcatg	attaagtagc	3660
tgctcaggaa	ttgcctttgg	ttacattgcc	taatggtttg	atgtcaattt	ttcttctgaa	3720
tctttatttt	agatcaaaat	ggagctacaa	aagtgttcaa	ctgtgtatgg	atttgctttt	3780
gttgactgtg	ctgccttgag	gttttgggtt	gggtccatca	gcgatgatgc	atcatgtgct	3840
gctcttggag	cgttattgat	gcaggtaagc	aagtgtattc	tgtatcttat	gtgtaccatg	3900
tgacttcctg	tgcatatatt	tgggttgcag	gaactaattc	tgaatcacca	tttggtatgt	3960
tttttccagg	tttctccaaa	ggaagtgtta	tatgacagta	aaggtaaact	gcttgtatcg	4020
ccagttgttt	tgttaaacag	aatttaaggt	aaatgacact	ggttaattta	aagtgcatac	4080
atgttgaaat	attgcagggc	tatcaagaga	agcacaaaág	gctctaagga	aatatacgtt	4140
gacaggtacc	atttcagtag	gcaagctaac	tgacaattta	accgctcacc	gaatgatagg	4200
tctcttaaac	attgctaatg	tagatgatgt	ttatgtttca	atctaatagg	gtctacggcg	4260
gtacagttgg	ctccagtacc	acaagtaatg	ggggatacag	atgctgctgg	agttagaaat	4320
ataatagaat	ctaacggata	ctttaaaggt	tcttctgaat	catggaactg	tgctgttgat	4380
ggtctaaatg	aatgtgatgt	tgcccttagt	gctcttggag	agctaattaa	tcatctgtct	4440
aggctaaagg	tgtgttggct	tgtttagttt	ttgcttttca	caaattaagc	aaaggaactt	4500
ttcataactt	acagtttcta	tctacttgca	gctagaagat	gtacttaagc	atggggatat	4560
ttttccatac	caagtttaca	ggggttgtct	cagaattgat	ggccagacga	tggtaaatct	4620
tgagatattt	aacaatagct	gtgatggtgg	tccttcaggc	aagtgcatat	ttcttttttg	4680
ataacttcaa	ctagagggca	gacatagaag	gaaaaattct	aatacttcgt	acggatctcc	4740
agtaagtaat	agccgatttt	tgtttaccta	tgtagggacc	ttgtacaaat	atcttgataa	4800
ctgtgttagt	ccaactggta	agcgactctt	aaggaattgg	atctgccatc	cactcaaaqa	4860
tgtagaaagc	atcaataaac	ggcttgatgt	agttgaagaa	ttcacggcaa	actcagaaag	4920

tatgcaaatc actggccagt atctccacaa acttccagac ttagaaagac tgctcggacg catcaagtot agogttogat catcagooto tgtgttgcot gotottotgg ggaaaaaagt 5040 5100 getgaaacaa egagtaagta teaateacaa gttttetgag taatgeette eatgagtagt ataggactaa aacattacgg gtctagctaa agactgttct ccttctttg caatgtctgg 5160 ttattcatta catttctctt aacttattgc attgcaggtt aaagcatttg ggcaaattgt 5220 gaaagggttc agaagtggaa ttgatctgtt gttggctcta cagaaggaat caaatatgat 5280 gagtttgctt tataaactct gtaaacttcc tatattagta ggaaaaagcg ggctagagtt 5340 5400 atttetttet caattegaag cagecataga tagegaettt ceaaattate aggtgeeeat ctatctttca tactttacaa caaaatgtct gtcactactc aaagcaatgc atatggctta 5460 gateteaact cacacceega ggateetaaa gggatttget tettatteet aatgtttttg 5520 5580 gatggtttga tttatttcta acttgaactt attaatcttg taccagaacc aagatgtgac agatgaaaac gctgaaactc tcacaatact tatcgaactt tttatcgaaa gagcaactca 5640 atggtctgag gtcattcaca ccataagctg cctagatgtc ctgagatctt ttgcaatcgc 5700 5760 agcaagtete tetgetggaa geatggeeag geetgttatt tetecegaat cagaagetae 5820 agatcagaat cagaaaacaa aagggccaat acttaaaatc caaggactat ggcatccatt 5880 tgcagttgca gccgatggtc aattgcctgt tccgaatgat atactccttg gcgaggctag 5940 aagaagcagt ggcagcattc atcctcggtc attgttactg acgggaccaa acatgggcgg aaaatcaact cttcttcgtg caacatgtct ggccgttatc tttgcccaag tttgtatact 6000 6060 cyttagataa ttactctatt ctttgcaatc agttcttcaa catgaataat aaattctgtt 6120 ttotgtotgo agottggotg ctacgtgoog tgtgagtott gogaaatoto cotogtggat 6180 actatcttca caaggettgg egeatetgat agaatcatga caggagagag taagttttgt totcaaaata ccaattooto gaactattta otcagatttt gtotgattgg acaaggtggt 6240 6300 tttgcttttt tttaggtacc tttttggtag aatgcactga gacagcgtca gttcttcaga 6360 atgcaactca ggattcacta gtaatccttg acgaactggg cagaggaact agtactttcg atggatacgc cattgcatac teggtaacct getettetee tteaacttat acttgttgat 6420 caacaaaaac atgcaattca ttttgctgaa acttattgat ttatatcagg tttttcgtca 6480 6540 cotggtagag aaagttcaat gtcggatgct ctttgcaaca cattaccacc ctctcaccaa

ggaattegeg teteaceeae gtgteacete gaaacacatg gettgegeat teaaatcaag 6600 atotgattat caaccacgtg gttgtgatca agacctagtg ttcttgtacc gtttaaccga 6660 gggagcttgt cctgagagct acggacttca agtggcactc atggctggaa taccaaacca 6720 agtggttgaa acagcatcag gtgctgctca agccatgaag agatcaattg gggaaaactt 6780 caagtcaagt gagctaagat ctgagttctc aagtctgcat gaagactggc tcaagtcatt 6840 6900 ggtgggtatt tetegagteg eccaeaacaa tgeececatt ggegaagatg actaegacae tttgttttgc ttatggcatg agatcaaatc ctcttactgt gttcccaaat aaatggctat 6960 gacataacac tatctgaagc tegttaagtc ttttgcttct ctgatgttta ttcctcttaa 7020 7080 aaaatgetta tatateaaaa aattgtttee tegattataa caagattata tatgtatetg teggtttage tatggtatat aatatatgta tgttcatgag attggtcaag agaaatacte 7140 7200 acaaacagta tattaagaag gaaatatgtt tatgcattaa tttaagtttc aagataaact gcaaataacc tcgactaaag ttgcaaagac caaacacaaa ttacaaaact tataagactt 7260 aagttotgaa ttooctaaaa ccaaaaaaaa aaacagaaca tattttgttg catotacaaa 7320 7380 caacacaaac ctacatagtt tataacttac tcatcactga gattaacatc agaatcattc tecatttett catetteact etcateatea teaceaceae catgatgatt etceteetet 7440 tcacgtaacc tagcaatctc actctgagct ctatcaacaa tctgcttctt ctgcaactcc 7500 aaatctctct gaaaatcagc tctcatcttc tccaactcct tcatttgctc tttcttactc 7560 7620 ttetecatet teteataaae etteceaaae eteteaaeag aateegeeaa eatettatae gaagcagcgt cattaacctt cttcctctcg tactcaacct catcatectc atcctcctcc 7680 tetteagaat caccaggaet atceateate teateaaace cattagaett atetaaataa 7740 7800 accttagtgt tcataaacac aaactcacct gaatcaacac cacaagctaa acctaaatcc 7860 gacttgggcg aaacacaaag caacatatcc aacttattga aaaacgacca tttacttgaa cctaaacctg atttctcaac cttaatcttc tettttctat acttcctctt caagtcatca 7920 atcattctcc tacattgcgt ctcagatttc tccatcctta gctcctcact cactttctca 7980 getaetteat tecaateete gtteeteaaa eteettetae eeaattgeaa aaacetatet 8040 8067 ccccaaactt caagcaacac aa